

Brief Report

Mortality in neonate bats infested by ectoparasite bugs in India; humans are at risk too

Shanti Lal Choubisa^{1,2,*}, Pallavi Choubisa³

¹ Department of Advanced Science and Technology, National Institute of Medical Science and Research, NIMS University Rajasthan, Jaipur 303121, Rajasthan, India

² Former Department of Zoology, Government Meera Girls College, Udaipur 303002, Rajasthan, India

³ Department of Obstetrics and Gynaecology, RNT Medical College and Pannadhay Janana Hospital, Udaipur 313001, Rajasthan, India

* **Corresponding author:** Shanti Lal Choubisa, choubisasl@yahoo.com

CITATION

Choubisa SL, Choubisa P. Mortality in neonate bats infested by ectoparasite bugs in India; humans are at risk too. *Environment and Public Health Research*. 2024; 2(2): 1778.
<https://doi.org/10.59400/ephr1778>

ARTICLE INFO

Received: 26 September 2024

Accepted: 19 November 2024

Available online: 12 December 2024

COPYRIGHT



Copyright © 2024 by author(s).

Environment and Public Health

Research is published by Academic

Publishing Pte. Ltd. This work is

licensed under the Creative

Commons Attribution (CC BY)

license.

<https://creativecommons.org/licenses/by/4.0/>

by/4.0/

Abstract: Human bed-bugs are well known and are found almost all over the world. Many types of bugs that infest various species of bats (Chiroptera: Mammalia) are found in different geographical areas or ecosystems, feeding on their blood. These small bugs, which are external parasitic insects, belong to the family Cimicidae of the order Hemiptera of the class Insecta of the phylum Arthropoda of the animal kingdom. Their bites can cause negative health effects in humans such as skin reactions, anemia, insomnia, anxiety, and panic attacks. Although their bites and bloodsucking do not kill any host. In general, bat-bugs infest adult and old bats. But, recently in Udaipur, Rajasthan, India, six neonates or neonate bats of small Asiatic yellow bats (*Scotophilus kuhlii*) were also found infested with bat-bugs ectoparasites belonging to the genus *Cimex* Linnaeus, 1758 (*Cimex pattoni*). Such a rare and unique instance has never been observed and reported earlier from any geographical area in the world. Interestingly, out of these, three were also found dead (66.6%), two were in a dying state, and one was in a critical condition or struggling for survival. Due to our limitation, the exact cause of death in these infested bat neonates with bug parasites has not been investigated. In the present communication, details of this case have been focused on and also focus on how much people are at risk of infestation from these external parasitic bat-bugs.

Keywords: Asian yellow bats (*Scotophilus kuhlii*); neonate bats; bat-bugs; ectoparasite; health; humans; infestation; insects; India

1. Introduction

There are many species of tiny bugs found all over the world that are external parasitic insects of the family Cimicidae, belonging to the order Hemiptera of the class Insecta of the phylum Arthropoda of the animal kingdom [1]. In India, these insects are commonly known as “Khatmal” (in Hindi). However, maximum study and information about them is limited to human bed-bugs (*Cimex lectularius*, *C. lectularius*, *C. hemipterus*, *C. rotundatus*, *C. piloselus*, *C. pipistrale*, etc.). Generally, these ectoparasite bugs prefer to live in cracks and crevices, and they can survive in an unfavorable environment without food (blood) for many months or even years [2]. These nocturnal insects are also well adapted anatomically, morphologically, and physiologically, due to which they can survive for a long time even in adverse environments and have evolved excellent organs to suck the blood of not only human beings but also of various species of bats (Chiroptera: Mammalia) living in deserted ruins, abandoned buildings, caves, houses, and cracks and hollows of trees

around human settlements. Interestingly, they do not spare even the small cliff swallow (*Petrochelidon pyrrhonota*) which lives in mud nests under houses, buildings, and bridges [3,4]. These insects also infest birds and have been reported in birds like chickens, owls, and woodpeckers. These insects are also called bed-bugs, bat-bugs, swallow-bugs, and poultry-bugs depending on their host species. But all insects of different host species, be it human beings, bats, or birds, look almost similar in morphology or structure, and their behavior is also almost similar. Bat-bugs are very similar to bed-bugs and are so similar in appearance that they are often mistaken for bed-bugs. Therefore, a microscopic examination is required to correctly identify them. Interestingly, these bat-bugs can also bite humans and cause negative health effects such as skin reactions, anemia, insomnia, anxiety, and panic attacks in people [5–7]. In general, bat-bugs infest adult and old bats but not neonate bats. But, recently in Udaipur, Rajasthan, India, neonate bats of small Asiatic yellow bats (*Scotophilus kuhlii*) (**Figure 1a–d**) were also found infested with bat-bug ectoparasites belonging to the genus *Cimex* Linnaeus, 1758 (*Cimex pattoni*) (**Figure 1d–f**). Such a rare and unique instance has never been observed and reported before from any geographical area in the world. Among these infested neonate bats, 66.66% are also found dead. In the present communication, details of this case have been discussed, and there is also a focus on how much people are at risk of infestation from these external parasitic bat-bugs.




Figure 1. Baby or neonate bats infested with ectoparasite bugs.

One of these neonate bats has died (a), while others infested with bugs are struggling to survive (b,c). Some bugs (d) have been isolated from the baby bats (c). Mature (e) and immature bugs (f) were recovered from the neonate bats (b,c). Arrow indicates bug parasite sucking blood at dorsal side of neck region.

2. Mortality in infested neonate bats with ectoparasite bugs

Recently, during the rainy season (July, 2024), six small babies of small Asian yellow bats (*S. kuhlii*) were found lying under a cement pipe protruding from the roof of a house in Udaipur city of Rajasthan (India). Of these, three were found dead, two were in a dying state, and one was in a critical condition or struggling for survival. On closer inspection, 3-4 small bugs were found stuck on the dorsal surface of all these neonates or babies. When these bat-bugs were pressed lightly, blood oozed out of them. In fact, all these bat-bugs were full of blood. Surprisingly, when they were separated from the neonate bats with the help of a thin wooden stick, some of them quickly ran back towards these babies and stuck to their bodies again. There were some who fell upside down during separation. But in a short while, they immediately got back upright and ran towards these babies and stuck to them again. Such aggressive behavior of theirs was never seen and reported before. How do these insects or bat-bugs recognize their host, and how are they attracted towards them? But in the absence of research, it is not right to say so.

In the present study we found 66.66% mortality in baby bats infested with bugs. It is well known that blood sucking by bugs can cause mild to severe anemia depending on the parasitism or parasitaemia [8–11]. Parasitic bugs also suck the blood of neonate bats and develop anemia, which can cause mortality in them. However, we have not investigated the hemoglobin (Hb) level in any infested neonate bats to confirm the evidence of anemia in them. On the other hand, it is also possible that they die due to starvation. However, without any concrete evidence, it is difficult to say what was the exact cause of death in these newborns, and it is not clear yet. The truth is that we have suddenly come across this case, or this case has come to us by chance. In fact, due to our limitations, we have not investigated to find out the cause of postnatal death, so this can be a subject of further research.

In the world, various species of bat-bug ectoparasites derived from different bat host species have been well studied and reported from different geographical provinces [12,13]. In India, the first report on bat-bugs was reported in 1912 [14]. Subsequently, more than 15 more species of bat-bugs belonging to the family Cimicidae have been reported in different species of bats surviving in or around human habitations, which have been well reviewed [15–23]. From the states of Rajasthan and Gujarat of the country, five species of bat-bugs (*Cimex ueshimai*, *C. pattoni*, *Cacodmus bharti*, *Leptocimex hiregaudari*, and *Stricticimex namru*) from six species of bats (*Pipistrellus dormeri dormeri*, *Tophozous longimanus*, *Scotophilus heathi heathi*, *Taphozous kachhensis kachhensis*, *Rhinopoma microphyllum microphyllum*, and *Tadarida aegyptiaca thomasi*) belonging to the Pteropidae family have also been reported [14,15]. Based on the literature, the current species of the ectoparasitic bug is *C. pattoni*, recovered from the pups of either *S. kuhlii* or *S. heathi heathi*. Though both species are small and have little difference in their

morphology, the habitats are quite different. However, the first species is more accurate and possible, as these small creatures also roost in hidden places (cracks, crevices, cemented pipes coming out of the roofs of houses, abandoned buildings, hollows of nearby trees, etc.) found in the vicinity of human habitation. The present neonate bats were also found in similar habitat or lying under a cement pipe coming out of the roof of a house. This species enters people's homes at night whenever it gets a chance. Which can also be a danger to humans. Because when these bats hide in houses, their bug parasitic insects also go to the houses along with them, which hide in the nearby cracks and bite the people of that house and suck their blood as soon as they get a chance. It is, generally, believed that the bite of these insects does not spread any infectious disease in humans. But human bed-bugs have been reported to be carriers of many pathogens [23–30]. Bed-bugs have also been found to be infected in nature with *Wuchereria bancrofti* and *Brugia malayi* (causes of filariasis), *Trypanosoma cruzi*, *Brucella melitensis*, *Coxiella burnetii*, and *Rickettsia*, which causes exanthematous typhus [13,23–30]. However, more authentic or scientific studies are still needed to confirm whether these bat parasites can cause and spread any infectious disease in humans.

3. Significance

The death of neonate bats suffering from bug parasites can affect their survival and population. The reduction in their population can harm the environment. Because bats are a natural source of controlling the population of various pests that harm economically important plants and agricultural crops [31,32]. Bats also disperse the seeds of fruit trees, thereby strengthening the existing ecosystem [33–35]. Due to the social behavior of bats and their proximity to human populations, bat-bug parasites can also be spread to humans. Much research is still needed in India to understand bat-bug parasitism and the behavior of different species of bats inhabiting human habitations. However, the significance of this report is that this is the first time that dead neonate or immature bats have been found infested with bug parasites, which is rare and unique. This important finding also adds to or expands the existing knowledge about parasitism in bats.

Acknowledgments: The authors are grateful to eminent zoologists A. N. Sharma and A. K. Purohit for their help in the identification of insects and bats and sharing related scientific knowledge, respectively. The authors also thank Darshana Choubisa, Department of Prosthodontics, Geetanjali Dental and Research Institute, Udaipur, Rajasthan 313002, India, for her support.

Conflict of interest: The authors declare no conflict of interest.

References

1. Kotpal RL. Modern Text Book of Zoology Invertebrates. Rastogi Publications; 2011.
2. Thompson L. Adult Bug Bites, Rashes and Symptoms. The University of Chicago; 2012.
3. Brown CR, Brown MB. Ectoparasitism as a cause of natal dispersal in Cliff Swallows. *Ecology*. 1992; 73(5): 1718-1723. doi: 10.2307/1940023

4. Brown CR, Brown MB. *Coloniality in the Cliff Swallow: The Effect of Group Size on Social Behavior*. University of Chicago Press, Chicago, IL; 1996.
5. Goddard J, de Shazo R. Rapid rise in bed bug populations: the need to include them in the differential diagnosis of mysterious skin rashes. *Southern Medical Journal*. 2008; 101(8): 854-855. doi: 10.1097/smj.0b013e31817be9db
6. Goddard J, de Shazo R. Bed bugs (*Cimex lectularius*) and clinical consequences of their bites. *Journal of American Medical Association*. 2009; 301(13): 1358-1366.
7. Goddard J, de Shazo R. Psychological Effects of Bed Bug Attacks (*Cimex lectularius* L.). *The American Journal of Medicine*. 2012; 125(1): 101-103. doi: 10.1016/j.amjmed.2011.08.010
8. Pritchard MJ, Hwang SW. Severe anemia from bedbugs. *Canadian Medical Association Journal*. 2009; 181(5): 287-288. doi: 10.1503/cmaj.090482
9. Paulke-Korinek M, Széll M, Laferl H, et al. Bed bugs can cause severe anaemia in adults. *Parasitology Research*. 2011; 110(6): 2577-2579. doi: 10.1007/s00436-011-2777-6
10. Sheele JM, Pritt BS, Libertin CR, et al. Bed bugs are associated with anemia. *The American Journal of Emergency Medicine*. 2021; 46: 482-488. doi: 10.1016/j.ajem.2020.10.070
11. Izri A, Marteau A, Ferreira T, et al. Severe anemia due to bed bugs hyperinfestation. *Microbial Pathogenesis*. 2020; 149: 104564. doi: 10.1016/j.micpath.2020.104564
12. Usinger RL. *Monograph of Cimicidae (Hemiptera: Heteroptera)*. Washington, D. C., The Thomas Say Foundation; 1996. pp. 585.
13. Ueshima N. New species and records of Cimicidae with keys (Hemiptera). *Pan Pacific Entomology*. 1968; 44 (4): 269-279.
14. Jordon K, Rothschild NC. Notes on Clinocoridae, a family of Rhynchota, with the descriptions of a new genus and species. *Novitates Zoologicae*. 1912; 19: 352-356.
15. Patton WS, Cragg FW. *A Textbook of Medical Entomology*. Published online 1913. doi: 10.5962/bhl.title.101563
16. Horváth G. Description of a new bat-bug from India. *Records of the Zoological Survey of India*. 1925; 27(3): 191. doi: 10.26515/rzsi/v27/i3/1925/163451
17. Mathur RN. A new species of bat-bug *Aphraniola* from Dehra Dun Cimicidae Hemiptera. *Indian Journal of Entomology*. 1953; 14: 257-261.
18. Hiregoudar LS, Bal DV. Some ectoparasites of bats from India. *Agra University of Journal Research (Science)*. 1956; 5: 1-134.
19. Bhat HR, Sreenivasan MA, Ilkal MA. *Cimex usingeri*, sp. nov. (hemiptera: cimicidae) from a colony of *Rhinolophus rouxii* Temminck, 1835 (chiroptera: rhinolophidae) in India. *Oriental Insects*. 1973; 7(1): 79-82. doi: 10.1080/00305316.1973.10434206
20. Bhat HR. *Cimex himalayanus*, sp. nov. (Hemiptera : Cimicidae) infesting a bat, *Myotis Siligorensis* in the Himalayan region of Uttar Pradesh, India. *Oriental Insects*. 1974; 8(2): 181-184. doi: 10.1080/00305316.1974.10434457
21. Bhat HR. A review of Indian Cimicidae (Hemiptera—Heteroptera). *Oriental Insects*. 1974; 8(4): 545-550. doi: 10.1080/00305316.1974.10434891
22. Advani R, Vazirani TG. Studies on ectoparasites of bats of Rajasthan and Gujarat (India). *Records of Zoological Survey of India*. 1981; 22: 1-155.
23. Burton GJ. Bedbugs in Relation to Transmission of Human Diseases: Review of the Literature. *Public Health Reports (1896-1970)*. 1963; 78(6): 513. doi: 10.2307/4591852
24. Blow JA, Turell MJ, Silverman AL, et al. Stercorarial shedding and transtadial transmission of hepatitis B virus by common bed bugs (Hemiptera: Cimicidae). *Journal of Medical Entomology*. 2001; 38(5): 694-700. doi: 10.1603/0022-2585-38.5.694
25. Goddard J, Varela-Stokes A, Smith W, et al. Artificial infection of the bed bug with *Rickettsia parkeri*. *Journal of Medical Entomology*. 2012; 49(4): 922-926. doi: 10.1603/me11282
26. Zorrilla-Vaca A. Bedbugs and vector-borne diseases. *Clinical Infectious Diseases*. 2014; 59(9): 1351-1352. doi: 10.1093/cid/ciu575
27. Barbarin AM, Hu B, Nachamkin I, et al. Colonization of *Cimex lectularius* with methicillin-resistant *Staphylococcus aureus*. *Environmental Microbiology*. 2014; 16(5): 1222-1224. doi: 10.1111/1462-2920.12384
28. Salazar R, Castillo-Neyra R, Tustin AW, et al. Bed bugs (*Cimex lectularius*) as vectors of *Trypanosoma cruzi*. *The American Society of Tropical Medicine and Hygiene*. 2015; 92(2): 331-335. doi: 10.4269/ajtmh.14-0483

29. Leulmi H, Bitam I, Berenger JM, et al. Correction: Competence of *Cimex lectularius* bed bugs for the transmission of *Bartonella quintana*, the agent of trench fever. PLOS Neglected Tropical Diseases. 2015; 9(6): e0003871. doi: 10.1371/journal.pntd.0003871
30. Lai O, Ho D, Glick S, et al. Bed bugs and possible transmission of human pathogens: a systematic review. Archives of Dermatological Research. 2016; 308(8): 531-538. doi: 10.1007/s00403-016-1661-8
31. Boyles JG, Cryan PM, McCracken GF, et al. Economic importance of bats in agriculture. Science. 2011; 332(6025): 41-42. doi: 10.1126/science.1201366
32. Boyles JG, Sole CL, Cryan PM, et al. On estimating the economic value of insectivorous bats: prospects and priorities for biologists. In: Bat Evolution, Ecology, and Conservation. Springer, New York, NY; 2013. pp. 501-515.
33. Whitby MD, Kieran TJ, Glenn TC, et al. Agricultural pests consumed by common bat species in the United States corn belt: The importance of DNA primer choice. Agriculture, Ecosystems & Environment. 2020; 303: 107105. doi: 10.1016/j.agee.2020.107105
34. Preciado-Benítez O, Gómez y Gómez B, Navarrete-Gutiérrez DA, et al. The use of commercial fruits as attraction agents may increase the seed dispersal by bats to degraded areas in Southern Mexico. Tropical Conservation Science. 2015; 8(2): 301-317. doi: 10.1177/194008291500800203
35. Mahandran V, Murugan CM, Marimuthu G, et al. Seed dispersal of a tropical deciduous Mahua tree, *Madhuca latifolia* (Sapotaceae) exhibiting bat-fruit syndrome by pteropodid bats. Global Ecology and Conservation. 2018; 14: e00396. doi: 10.1016/j.gecco.2018.e00396