

# On the relation between perfect powers and tetration frozen digits

Marco Ripà

Independent Researcher, Rome, Italy; marcokrt1984@yahoo.it

## CITATION

Ripà M. On the relation between perfect powers and tetration frozen digits. Journal of AppliedMath. 2024; 2(5): 1771.  
<https://doi.org/10.59400/jam1771>

## ARTICLE INFO

Received: 25 September 2024  
Accepted: 4 November 2024  
Available online: 9 December 2024

## COPYRIGHT



Copyright © 2024 author(s).  
Journal of AppliedMath is published by Academic Publishing Pte. Ltd. This work is licensed under the Creative Commons Attribution (CC BY) license.  
<https://creativecommons.org/licenses/by/4.0/>

**Abstract:** This paper provides a link between integer exponentiation and integer tetration since it is devoted to introducing some peculiar sets of perfect powers characterized by any given value of their constant congruence speed, revealing a fascinating relation between the degree of every perfect power belonging to any congruence class modulo 20 and the number of digits frozen by these special tetration bases, in radix-10, for any unit increment of the hyperexponent. In particular, given any positive integer  $c$ , we constructively prove the existence of infinitely many  $c$ -th perfect powers having a constant congruence speed of  $c$ .

**Keywords:** congruence speed; convergence; perfect power; power tower; tetration

## 1. Introduction

We explore a tool in modular arithmetic that predicts patterns among the right-hand digits of terms in rapidly growing sequences arising from power towers, thereby reducing computational overhead. One primary application of this approach is in cryptography, particularly within modular exponentiation algorithms. With the possible advent of quantum computers, this method could potentially improve the performance and security of encryption algorithms. Additionally, the results discussed here have implications for enhancing the efficiency of pseudorandom number generators (PRNGs) that rely on modular arithmetic.

For clarity, we will denote  $\mathbb{N}_0$  as the set of nonnegative integers (including zero) and  $\mathbb{N}$  as the set of positive integers  $\{1, 2, 3, \dots\}$ .

In recent years, by assuming the standard decimal numeral system (radix-10), we have shown that the integer tetration  ${}^b a := \begin{cases} a & \text{if } b = 1 \\ a({}^{b-1} a) & \text{if } b \geq 2 \end{cases}$  has a unique property [1] involving the number of new frozen rightmost digits for any unit increment of its hyperexponent,  $b \in \mathbb{N}$  [2,3]. Indeed, this value no longer depends on  $b$  as  $b$  becomes sufficiently large (see the sequence A372490 in the On-Line Encyclopedia of Integer Sequences [4]) and the tetration base,  $a \in \mathbb{N}$ , is not a multiple of 10.

We refer to the mentioned property as the constancy of the congruence speed of tetration (see **Definition 1** and also the comments of the OEIS sequence A317905).

From [2], we know that each positive integer  $a \geq 2$ , which is not a multiple of 10, is characterized by a finite, strictly positive, integer value of its constant congruence speed (the map of the constant congruence speed of every  $a$  as above is provided by [2,3]).

Then, the present paper aims to constructively prove the existence of infinitely many perfect powers having any given positive constant congruence speed.

A pleasant result, that follows from **Theorem 3** as a corollary, is the existence, for any given positive integer  $c$ , of infinitely many  $c$ -th perfect powers (i.e., an integer  $a > 1$  is a  $c$ -th perfect power if there exist some integers  $\tilde{a}$  and  $c$  such that

$a = \tilde{a}^c$ , so we have perfect squares if  $c = 2$ , perfect cubes if  $c = 3$ , and so forth) having a constant congruence speed of  $c$ .

## 2. Preliminary investigations with the automorphic numbers

In order to present the results compactly, let us properly define the constant congruence speed of tetration as already done in [3, p. 442], Definitions 1.1 and 1.2.

**Definition 1.** Let  $n \in \mathbb{N}_0$  and assume that  $a \in \mathbb{N} - \{1\}$  is not a multiple of 10. Then, given  ${}^{b-1}a \equiv {}^b a \pmod{10^n} \wedge {}^{b-1}a \not\equiv {}^b a \pmod{10^{n+1}}$ , for all  $b \in \mathbb{N}$ ,  $V(a, b)$  returns the nonnegative integer such that  ${}^b a \equiv {}^{b+1} a \pmod{10^{n+V(a,b)}} \wedge {}^b a \not\equiv {}^{b+1} a \pmod{10^{n+V(a,b)+1}}$ , and we define  $V(a, b)$  as the congruence speed of the base  $a$  at the given height of its hyperexponent  $b$ .

Furthermore, let  $\bar{b} := \min\{b \in \mathbb{N} : V(a, b) = V(a, b + k) \text{ for all } k \in \mathbb{N}\}$  so that we define, as constant congruence speed of  $a$ , the positive integer  $V(a) := V(a, \bar{b})$ .

In general, we know that a sufficient but not necessary condition for having  $V(a) = V(a, \bar{b})$  is to set  $\bar{b} := a + 1$ , and for a tighter bound on  $\bar{b} := \bar{b}(a)$ , holding for any  $a \not\equiv 0 \pmod{10}$ , see [3, p. 450].

As a clarifying example, let us consider the case  $a = 807$ . Then, we have  $V(807, 1) = 0$ ,  $V(807, 2) = 4$ ,  $V(807, 3) = 4$ ,  $V(807, 4) = 4$ ,  $V(807, 5) = 4$ , and finally  $V(807, 6) = V(807, 7) = \dots = V(807) = 3$  since  $\tilde{v}_5(807^2 + 1) + 2 = 4 + 2 = 6$  (by [3, Definition 2.1]) and

$$\begin{aligned} {}^1_{807} &= 807; \\ {}^2_{807} &\equiv 549620396283318273888501737943 \pmod{10^{30}}; \\ {}^3_{807} &\equiv 601692651466822940525632857943 \pmod{10^{30}}; \\ {}^4_{807} &\equiv 146336906474874632626032857943 \pmod{10^{30}}; \\ {}^5_{807} &\equiv 355034907448973150626032857943 \pmod{10^{30}}; \\ {}^6_{807} &\equiv 478635689812283150626032857943 \pmod{10^{30}}; \\ {}^7_{807} &\equiv 027048888762283150626032857943 \pmod{10^{30}}; \\ &\vdots \\ {}^{(1 \text{ googoleplex})}_{807} &\equiv 803001638762283150626032857943 \pmod{10^{30}}. \end{aligned}$$

**Lemma 1.** Let  $a \in \mathbb{N}$  be such that  $a \not\equiv 0 \pmod{10}$ . Then, for all  $t \in \mathbb{N}_0$ , there exist infinitely many  $c \in \mathbb{N}$  such that  $V(a^c) = t$ .

**Proof of Lemma 1.** Disregarding the special case  $t = 0$ , this proof immediately follows from **Definition 1**.  $\square$

For any integer  $a > 1$  which is not a multiple of 10, the constant congruence speed of the tetration  ${}^b a$  is well-defined and it is the same for any  $b \in \{a + 1, a + 2, a + 3, \dots\}$ . Thus, by the last line of Equation (2) in [2], it is sufficient to consider  $\hat{a} := 10^t - 1$  so that  $V(\hat{a}) = t$  is true for any given positive integer  $t$ , and

then we can easily complete the proof by observing that  $V(1) = 0$  as stated in [3, Definition 1.3].

Trivially,  $V(\hat{a}, b) = V(\hat{a}, b + 1) = V(\hat{a}, b + 2) = \dots$  is certainly true for every integer  $b \geq \hat{a} + 1$  and, by assuming that  $t \in \mathbb{N}$ , for all the aforementioned values of  $b$ , we have that  $\hat{a} = 10^t - 1$  implies  $V(\hat{a}, b) = t$  (while from  $a = 1 \Rightarrow t = 0$  it follows that  $V(1^c) = 0$  for any nonnegative integer  $c$ ).

Consequently, let  $t \in \mathbb{N}$ , assume  $b \in \{10^t, 10^t + 1, 10^t + 2, \dots\}$ , and then  $V((10^t - 1)^c, b) = V((10^t - 1)^c) = t$  is true for any  $c \in \{b^{-1}(10^t - 1), b(10^t - 1), b^{+1}(10^t - 1), \dots\}$  so that the **proof of Lemma 1** is complete.

Thus, **Lemma 1** shows the existence of infinitely many  $c$ -th powers of the tetration base  $a : a \equiv 1, 2, 3, 4, 5, 6, 7, 8, 9 \pmod{10}$  that are characterized by any given (arbitrarily large) nonnegative constant congruence speed.

**Remark 1.** We note that, in radix-10, there exist only three positive 1-automorphic numbers and they are congruent modulo 100 to 1, 25, and 76 (respectively). Thus, the corresponding three integers found by considering the two rightmost digits of the analogous solutions of the fundamental 10-adic (decadic) equation  $y^5 = y$ , by [2, Equation (2)] (see also the OEIS sequences A018247 and A018248), describe 1-automorphic numbers [5] (e.g.,  $\alpha_{76} \mapsto a_{76} := 76$  since  $76^2 \equiv 76 \pmod{10^2}$ ) and we know that [6], in radix-10, there are only four 10-adic solutions, including  $\alpha_{00} := \dots 000000$ , to the equation  $y^2 = y$ . Consequently, by looking at lines 4, 5, and 7 of Equation (16) in [3], we can see that the recurrences described by Equations (1)–(3) hold for every  $c \in \mathbb{N}$ .

$$\tilde{a} \equiv 6 \pmod{10} \Rightarrow V(\tilde{a}) \begin{cases} = V(\tilde{a}^c) \text{ iff } c \equiv 1, 2, 3, 4 \pmod{5} \\ \leq V(\tilde{a}^c) \text{ iff } c \equiv 0 \pmod{5} \end{cases} \quad (1)$$

$$\tilde{a} \equiv 5 \pmod{20} \Rightarrow V(\tilde{a}) \begin{cases} = V(\tilde{a}^c) \text{ iff } c \equiv 1 \pmod{2} \\ \leq V(\tilde{a}^c) \text{ iff } c \equiv 0 \pmod{2} \end{cases} \quad (2)$$

$$\tilde{a} \equiv 1 \pmod{20} \Rightarrow V(\tilde{a}) \begin{cases} = V(\tilde{a}^c) \text{ iff } c \equiv 1, 2, 3, 4 \pmod{5} \\ \leq V(\tilde{a}^c) \text{ iff } c \equiv 0 \pmod{5} \end{cases} \quad (3)$$

The investigation of this observation (with specific reference to Equation (2)) leads us to the following theorem.

**Theorem 1.** For each  $c \in \mathbb{N}$ , there exist infinitely many  $a : a \equiv 5 \pmod{20}$  such that  $\sqrt[c]{a} \in \mathbb{N} \wedge V(\sqrt[c]{a}) = t \wedge V(a) \geq t$  holds for all  $t \in \mathbb{N} - \{1\}$ . Symmetrically, for each  $t \in \mathbb{N} - \{1\}$ , there exist infinitely many  $a : a \equiv 5 \pmod{20}$  such that  $\sqrt[c]{a} \in \mathbb{N} \wedge V(\sqrt[c]{a}) = t \wedge V(a) \geq t$  holds for all  $c \in \mathbb{N}$ .

**Proof of Theorem 1.** Let us (constructively) prove first the last statement of **Theorem 1** since it simply follows from the constancy of the congruence speed as it has been shown in [2, Section 2.1].  $\square$

Let the symbol “\_” indicate the juxtaposition of consecutive digits (e.g.,  $3\_6\_1 = 36\_1 = 3\_61 = 361$ ). Consider the rightmost  $t \in \mathbb{N} - \{1\}$  digits of the 10-adic integer  $\alpha_{25} := \{5^{2^n}\}_\infty$ , say  $x_t\_x_{t-1}\_ \dots \_2\_5$ , and then juxtapose to the left the  $(t + 1)$ -th digit plus 1 if  $x_{t+1} \leq 8$  or the  $(t + 1)$ -th digit minus 1 if  $x_{t+1} = 9$ .

$$\text{So, let } \tilde{x}_{t+1} := \begin{cases} x_{t+1} + 1 & \text{if } x_{t+1} \in \{0, 1, 2, 3, 4, 5, 6, 7, 8\} \\ x_{t+1} - 1 & \text{if } x_{t+1} \in \{9\} \end{cases}$$

Thus, the base  $\tilde{a} := \tilde{x}_{t+1}x_t x_{t-1} \dots 2_5$  is characterized by a constant congruence speed of  $t$  (i.e.,  $V(\tilde{x}_{t+1}x_t x_{t-1} \dots 2_5) = V(\tilde{a}) = t$  for any  $t \in \mathbb{N} - \{1\}$ ). This property follows from [3, Equation (16)] (i.e.,  $2^t \mid \tilde{x}_{t+1}x_t x_{t-1} \dots 2_5 \wedge 2^{t+1} \nmid \tilde{x}_{t+1}x_t x_{t-1} \dots 2_5$ , for any  $t \geq 2$ ). Since (as discussed in **Remark 1**)  $\alpha_{25} \mapsto a_{25} := 25$  and  $25^2 \equiv 25 \pmod{10^2}$ , from Hensel's lemma [7] (see also [8,9]), we have that if  $\tilde{a} \equiv \alpha_{25} \pmod{10^t} \wedge \tilde{a} \not\equiv \alpha_{25} \pmod{10^{t+1}}$ , then  $\tilde{a}^c \equiv \alpha_{25} \pmod{10^t}$  for any given  $c \in \mathbb{N}$  (in general, we cannot assert that  $(\tilde{a} \equiv \alpha_{25} \pmod{10^t} \wedge \tilde{a} \not\equiv \alpha_{25} \pmod{10^{t+1}})$  implies  $\tilde{a}^c \not\equiv \alpha_{25} \pmod{10^{t+1}}$  for the given pair  $(t, c)$ ).

Consequently, by simply taking  $a := (\tilde{x}_{t+1}x_t x_{t-1} \dots 2_5)^c$  (as  $c$  is free to run over the positive integers), we have proven the existence, for any given  $t \in \mathbb{N} - \{1\}$ , of infinitely many tetration bases  $a \equiv 5 \pmod{20}$  such that  $V(a) \geq t$  holds for all the elements of the aforementioned set, a set that contains infinitely many distinct perfect powers originated from the string  $\tilde{x}_{t+1}x_t x_{t-1} \dots 2_5$  (since  $\tilde{a}$  is a positive integer by definition, then  $a = \tilde{a}^c$  implies that  $\sqrt[c]{a} \in \mathbb{N}$ ). Hence,  $V(\tilde{x}_{t+1}x_t x_{t-1} \dots 2_5) = t$  implies  $V((\tilde{x}_{t+1}x_t x_{t-1} \dots 2_5)^c) \geq t$  by observing that  $V((\tilde{x}_{t+1}x_t x_{t-1} \dots 2_5)^c) = V(\dots x_t x_{t-1} \dots 2_5)$ , and trivially  $\sqrt[c]{(\tilde{x}_{t+1}x_t x_{t-1} \dots 2_5)^c} \in \mathbb{N}$  for all  $c \in \mathbb{N}$  (we point out that, for any  $t \geq 2$  and as long as  $c$  is a positive integer,  $(\tilde{x}_{t+1}x_t x_{t-1} \dots 2_5)^c \equiv x_t x_{t-1} \dots 2_5 \pmod{10^t}$  holds by construction [5]).

Now, let us prove the first statement of **Theorem 1** and complete the proof.

For this purpose, it is sufficient to note that

$$V(\tilde{x}_{t+1}x_t x_{t-1} \dots 2_5) = V(10^{k+t} + \tilde{x}_{t+1}x_t x_{t-1} \dots 2_5)$$

is true for any positive integer  $k$ . So, we can take the  $c$ -th power of every integer of the form  $10^{k+t} + \tilde{x}_{t+1}x_t x_{t-1} \dots 2_5$  to get  $k$  distinct sets of cardinality  $\aleph_0$  each, whose elements, by construction, always satisfy the first statement of the theorem (we have already shown that, for any given  $c \in \mathbb{N}$ , if  $V(10^{k+t} + \tilde{x}_{t+1}x_t x_{t-1} \dots 2_5) = t$ , then  $V((10^{k+t} + \tilde{x}_{t+1}x_t x_{t-1} \dots 2_5)^c) \geq t$  holds for every  $t \in \mathbb{N} - \{1\}$ ).

Therefore, both statements of **Theorem 1** have been shown to be true, and this concludes the proof.

### 3. Main result

From here on, let us indicate the  $p$ -adic valuation [10] of any tetration base  $a$  as  $v_p(a)$ , for any prime number  $p$ .

Then, we need the following lemma to prove the existence, for any  $\tilde{a} \in \mathbb{N} - \{1\}$  such that  $\tilde{a} \not\equiv 0 \pmod{10}$ , of infinitely many  $c$ -th powers of  $\tilde{a}$  having a constant congruence speed of  $V(\tilde{a})$ ,  $V(\tilde{a}) + 1$ ,  $V(\tilde{a}) + 2$ ,  $V(\tilde{a}) + 3$ , and so forth. Furthermore, for any given positive integer  $c$ , **Lemma 2** (thanks to [3], Equation (2), line 2) implies the existence of infinitely many tetration bases of the form

$(10^{k+t} + 10^t + 1)^c$  (where  $k \in \mathbb{N}$  and  $t \in \mathbb{N} - \{1\}$ ) characterized by any constant congruence speed greater than  $1 + \min\{v_5(c), v_2(c)\}$ .

**Lemma 2.** For every  $c, k \in \mathbb{N}$  and  $t \in \mathbb{N} - \{1\}$ ,  $v_5\left(\left(10^{k+t} + 10^t + 1\right)^c - 1\right) = t + v_5(c)$  and  $v_2\left(\left(10^{k+t} + 10^t + 1\right)^c - 1\right) = t + v_2(c)$ .

**Proof of Lemma 2.** First of all, we prove that, for every positive integer  $c$ ,  $v_5\left(\left(10^{k+t} + 10^t + 1\right)^c - 1\right) = t + v_5(c)$ , where  $t, k \in \mathbb{N}$ .  $\square$

This can be achieved by using the lifting-the-exponent lemma (LTE lemma).

For this purpose, let  $c$  be a positive integer, we note that

$$v_5\left(\left(10^{k+t} + 10^t + 1\right)^c - 1\right) = v_5\left(\left(10^{k+t} + 10^t + 1\right)^c - 1^c\right),$$

so we can invoke the LTE lemma (see [11] and [12, Lemma 2.6]) for odd primes, stating that for any integers  $x, y$ , a positive integer  $c$ , and a prime number  $p$  such that  $p \nmid x \wedge p \nmid y$ , if  $p$  divides  $x - y$ , then  $v_p(x^c - y^c) = v_p(x - y) + v_p(c)$ .

Thus, by observing that  $p := 5$  is an odd prime satisfying all the conditions above for  $x := 10^{k+t} + 10^t + 1 \wedge y := 1$  (since  $10^{k+t} + 10^t$  is a multiple of 5),

$$v_5\left(\left(10^{k+t} + 10^t + 1\right)^c - 1^c\right) = v_5(10^{k+t} + 10^t) + v_5(c) = t + v_5(c).$$

Similarly, we can use the LTE lemma to show that  $2|c \Rightarrow v_2\left(\left(10^{k+t} + 10^t + 1\right)^c - 1\right) = t + v_2(c)$  and also  $2 \nmid c \Rightarrow v_2\left(\left(10^{k+t} + 10^t + 1\right)^c - 1\right) = t + v_2(c)$ .

In detail (see [13]), both  $x := 10^{k+t} + 10^t + 1$  and  $y := 1$  are odd numbers so that we can invoke the LTE lemma for  $p = 2$  since  $2 \nmid x$ ,  $2 \nmid y$ , and  $2|(x - y)$ .

Thus, the LTE lemma for  $p = 2$  states that if  $c$  is odd, then  $v_2\left(\left(10^{k+t} + 10^t + 1\right)^c - 1^c\right) = v_2(10^{k+t} + 10^t + 1 - 1) = v_2(10^t) = t$ . Since  $2 \nmid c \Rightarrow v_2(c) = 0$ , we can safely rewrite the above as  $v_2\left(\left(10^{k+t} + 10^t + 1\right)^c - 1\right) = t + v_2(c)$ .

On the other hand, if  $c$  is even, from the LTE lemma it follows that

$$\begin{aligned} & v_2\left(\left(10^{k+t} + 10^t + 1\right)^c - 1^c\right) \\ &= v_2(10^{k+t} + 10^t + 1 - 1) + v_2(10^{k+t} + 10^t + 1 + 1) + v_2(c) - 1 \\ &= v_2(10^{k+t} + 10^t) + v_2(10^{k+t} + 10^t + 2) + v_2(c) - 1. \end{aligned}$$

Now, for any given pair of positive integers  $(r, s)$  and a prime  $p$ , we know that  $v_p(r + s) = \min\{v_p(r), v_p(s)\}$  as long as  $v_p(r) \neq v_p(s)$ . Accordingly,  $t \geq 2$  implies  $v_2(10^t) \geq 2 > v_2(2)$  so that  $v_2(10^{k+t} + 10^t + 2) = \min\{v_2(10^{k+t}), v_2(10^t), v_2(2)\} = 1$ .

Hence,  $2|c \Rightarrow v_2\left(\left(10^{k+t} + 10^t + 1\right)^c - 1\right) = t + 1 - 1 + v_2(c) = t + v_2(c)$ .

Therefore, for every triad  $(t, c, k)$  of positive integers,  $v_5\left(\left(10^{k+t} + 10^t + 1\right)^c - 1\right) = t + v_5(c)$  and, symmetrically,  $v_2\left(\left(10^{k+t} + 10^t + 1\right)^c - 1\right) = t + v_2(c)$ . This completes the proof.

**Theorem 2.** For any given triad  $c, k \in \mathbb{N}$  and  $t \in \mathbb{N} - \{1\}$ ,

$$V\left((10^{k+t} + 10^t + 1)^c\right) = t + \min\{v_5(c), v_2(c)\}.$$

**Proof of Theorem 2.** **Theorem 2** easily follows from **Lemma 2**. Since  $t \geq 2$ ,  $10^{k+t} + 10^t + 1 \equiv 1 \pmod{10^2}$  (the congruence modulo 100 does not depend on  $k$  as  $k + t > t$ ), and then also  $(10^{k+t} + 10^t + 1)^c \equiv 1 \pmod{100}$  holds for each positive integer  $c$ . By **Theorem 2.1** of [3] (see Equation (2), line 2), we have that  $V\left((10^{k+t} + 10^t + 1)^c\right) = \min\left\{v_5\left((10^{k+t} + 10^t + 1)^c - 1\right), v_2\left((10^{k+t} + 10^t + 1)^c - 1\right)\right\}$ . Since **Lemma 2** asserts that  $v_5\left((10^{k+t} + 10^t + 1)^c - 1\right) = t + v_5(c)$  and  $v_2\left((10^{k+t} + 10^t + 1)^c - 1\right) = t + v_2(c)$ , it follows that  $V\left((10^{k+t} + 10^t + 1)^c\right) = \min\{t + v_5(c), t + v_2(c)\} = t + \min\{v_5(c), v_2(c)\}$  for any positive integers  $c, k$ , and  $t - 1$ .  $\square$

**Remark 2.** Let the tetration base  $a : a \equiv 6 \pmod{10}$  be given. Then, by looking at the two rightmost digits of the corresponding 10-adic solution,  $\alpha_{76} = \dots 7109376$  (see **Remark 1**), and applying the usual strategy (already described in the **proof of Theorem 1**), we find the sequence  $a_n := 10^{n+1} + 86, n \in \mathbb{N}$  (defining also the set  $\{186, 1086, 10086, 100086, \dots\}$ ). Now, we can create an infinite set consisting of the  $c$ -th powers of each aforementioned term, a set whose elements are all characterized by a unit constant congruence speed as long as  $c$  is not a multiple of 5. Thus,  $5 \nmid c$  implies  $V(186^c) = V(1086^c) = V(10086^c) = V(100086^c) = \dots = 1$ .

The above is just another example of the  $\tilde{x}_{t+1}$  idea, introduced in the **proof of Theorem 1**, shown by taking into account  $t = 1$  and the solution  $\alpha_{76} := 1 - \{5^{2^n}\}_\infty$  of the equation  $y^2 = y$  in the commutative ring of 10-adic integers (as we know [2], the other three solutions are  $\alpha_{00} := 0, \alpha_{01} := 1$ , and  $\alpha_{25} := \{5^{2^n}\}_\infty$ ).

**Corollary 1.** Let  $t \in \mathbb{N}$  and assume that  $V(\tilde{a}) = t$ . Then, for any nonnegative integer  $h$ , there exist infinitely many  $c \in \mathbb{N}$  such that  $V(\tilde{a}^c) = t + h$ .

**Proof of Corollary 1.** Let  $k \in \mathbb{N}$ . If  $t > 1$ , it is sufficient to observe that, by **Theorem 2**,

$$v_2(c) \geq v_5(c) \Rightarrow V\left((10^{k+t} + 10^t + 1)^c\right) = t + v_5(c).$$

Accordingly, let  $\tilde{a} := 10^{k+t} + 10^t + 1, c := 2^{k+h} \cdot 5^h$  (so that  $v_5(2^{k+h}) \leq v_2(2^{k+h})$ ), and then, for any  $h \in \mathbb{N}_0, \tilde{a}^c$  identifies an infinite set of valid tetration bases (since  $v_5(2^{k+h} \cdot 5^h) = h$  is true for any positive integer  $k$  and, consequently, the constant congruence speed of  $\tilde{a}^{2^{k+h} \cdot 5^h}$  does not depend on  $k$ ).  $\square$

For the remaining case,  $t = 1$ , **Remark 2** gives us a valid set of solutions (i.e., any tetration base of the form  $(10^{k+1} + 86)^{2^{k-1} \cdot 5^h}$  does the job since  $V(10^{k+1} + 86) = 1$  for every positive integer  $k$ ), so just let  $\tilde{a} := 10^{k+1} + 86$  and  $c := 2^{k-1} \cdot 5^h$  in order to get  $V\left((10^{k+1} + 86)^{2^{k-1} \cdot 5^h}\right) = v_5\left((10^{k+1} + 86)^{2^{k-1} \cdot 5^h} - 1\right) = 1 + v_5(5^h) = 1 + h$  (by [3], Equation (16), line 4).

**Theorem 3.** Let the integer  $a > 1$  not be a multiple of 10. Then, for all  $c \in \mathbb{N}$ , there exist infinitely many  $a$  such that  $\sqrt[c]{a} \in \mathbb{N} \wedge V(a) = V(\sqrt[c]{a}) = t$  holds for every integer  $t$  greater than  $1 + \min\{v_5(c), v_2(c)\}$ .

**Proof of Theorem 3.** Since **Theorem 2** states that, given  $t \in \mathbb{N} - \{1\}$ ,  $V\left((10^{k+t} + 10^t + 1)^c\right) = t + \min\{v_5(c), v_2(c)\}$  for each  $c, k \in \mathbb{N}$ , it follows that, for any given triad  $(t, k, c)$  such that  $t \geq 2 + \min\{v_5(c), v_2(c)\}$ , at least one of the two tetration bases  $(10^{k+t-v_5(c)} + 10^{t-v_5(c)} + 1)^c$  and  $(10^{k+t-v_2(c)} + 10^{t-v_2(c)} + 1)^c$  is guaranteed to have a constant congruence speed of  $t$ .  $\square$

Although this is enough to prove the theorem, we are free to simplify the generic form of the above by observing that (see [13]), as we call  $j := j(c)$  the last nonzero digit of  $c$ , only the following two cases matter:  $j \neq 5$  and  $j = 5$ .

Hence,

$$\begin{aligned} j \neq 5 &\Rightarrow v_5(c) \leq v_2(c) \\ \Rightarrow V\left((10^{k+t} + 10^{t-v_5(c)} + 1)^c\right) &= v_5\left((10^{k+t} + 10^{t-v_5(c)} + 1)^c - 1^c\right) \\ &= v_5(10^{k+t} + 10^{t-v_5(c)} + v_5(c)) \\ &= \min\{v_5(10^{k+t}), v_5(10^{t-v_5(c)})\} + v_5(c) \end{aligned}$$

(since  $k > -v_5(c)$  implies that  $v_5(10^{k+t}) > v_5(10^{t-v_5(c)})$ ), and then it follows that

$$V\left((10^{k+t} + 10^{t-v_5(c)} + 1)^c\right) = v_5(10^{t-v_5(c)} + v_5(c)) = t - v_5(c) + v_5(c) = t.$$

Conversely (given the fact that  $v_2(10^{k+t}) > v_2(10^{k+t-v_2(c)}) > v_2(10^{t-v_2(c)})$ ),

$$\begin{aligned} j = 5 &\Rightarrow v_5(c) > v_2(c) \\ \Rightarrow V\left((10^{k+t} + 10^{t-v_2(c)} + 1)^c\right) &= v_2\left((10^{k+t} + 10^{t-v_2(c)} + 1)^c - 1^c\right) \\ &= v_2\left((10^{k+t-v_2(c)} + 10^{t-v_2(c)} + 1)^c - 1\right) \end{aligned}$$

and, by **Lemma 2**,  $v_2\left((10^{k+t-v_2(c)} + 10^{t-v_2(c)} + 1)^c - 1\right) = t$  follows.

Thus,  $c : j \neq 5$  guarantees that  $V\left((10^{k+t} + 10^{t-v_5(c)} + 1)^c\right) = t$  is true for any  $c, k \in \mathbb{N}$ , and  $t > v_5(c)$ , while  $V\left((10^{k+t} + 10^{t-v_2(c)} + 1)^c\right) = t$  as  $c : j = 5$  and  $t > v_2(c)$ .

Therefore, we have shown that, for any given positive integer  $c$ , at least one of  $V\left((10^{k+t} + 10^{t-v_5(c)} + 1)^c\right) = t$  ( $t > v_5(c) + 1$ ) and  $V\left((10^{k+t} + 10^{t-v_2(c)} + 1)^c\right) = t$  ( $t > v_2(c) + 1$ ) is true, so the proof is complete.

Now, let  $t = c$  and observe that  $c - v_5(c) < 2 \Rightarrow c = 1$ , while  $c - v_2(c) < 2 \Rightarrow c \leq 2$ .

Then, the **proof of Theorem 3** shows the existence of a very special set of tetration bases that are  $c$ -th powers of an integer and whose constant congruence speed is  $c \in \mathbb{N} - \{1, 2\}$ , a set including all the bases of the form  $(10^{c+k} +$

$10^{c-\min\{v_5(c), v_2(c)\}} + 1)^c$  ,  $k \in \mathbb{N}$  (e.g.,  $(c = 1000 \wedge k = 314)$  ) implies  $V((10^{1314} + 10^{997} + 1)^{1000}) = 1000$ .

Therefore, for each  $c > 2$ , we have already proved **Corollary 2** as a particular case of **Theorem 3**.

**Corollary 2.** *Let  $a \in \mathbb{N} - \{1\}$  not be a multiple of 10. Then, for every  $c \in \mathbb{N}$ , there exist infinitely many  $a$  such that  $\sqrt[c]{a} \in \mathbb{N} \wedge V(a) = V(\sqrt[c]{a}) = c$ .*

Indeed, setting  $S$  as the sum of digits function to base 10 and assuming  $c > 2$ , we have  $S(10^{c+k} + 10^{c-\min\{v_5(c), v_2(c)\}} + 1) = 3$ ,  $3|10^{c+k} + 10^{c-\min\{v_5(c), v_2(c)\}} + 1$  and  $3^2 \nmid 10^{c+k} + 10^{c-\min\{v_5(c), v_2(c)\}} + 1$  (implying that none of these numbers can be a perfect power of degree greater than 1), so there exist infinitely many perfect powers of degree  $c = 3, 4, 5, 6, \dots$  having a constant congruence speed of  $c$ , and it is sufficient to consider all the tetration bases of the form  $(10^{c+k} + 10^{c-\min\{v_5(c), v_2(c)\}} + 1)^c$ .

On the other hand, if  $c \in \{1, 2\}$ , then the second last nonzero digit of any tetration base of the form  $(10^{c+k} + 10^{c-v_5(c)} + 1)^c$  will not be equal to 5 (we note that the second last nonzero digit of each  $c$ -th perfect power of  $10^{k+t} + 10^t + 1$  is always equal to the last nonzero digit of  $c$ ), and thus we can cover the remaining cases  $c = 1$  and  $c = 2$  by taking  $10^{1+k} + 10^1 + 1$  (since  $V((10^{1+k} + 11)^1) = 1$  holds for each positive integer  $k$  by [3], Equation (16), line 9) and  $(10^{2+k} + 10^2 + 1)^2$  (respectively).

Since  $\sqrt[c]{10^{1+k} + 11} \in \mathbb{N} \Rightarrow c = 1$  and  $\sqrt[c]{(10^{2+k} + 101)^2} \in \mathbb{N} \Rightarrow c \in \{1, 2\}$ , we have finally proved for any given positive integer  $c$  the existence of infinitely many tetration bases  $\tilde{a}^c$  which are perfect powers of degree  $c$  (exactly) and such that  $V(\tilde{a}^c) = V(\tilde{a}) = c$ .

**Remark 3.** *Let  $c, k \in \mathbb{N}$ ,  $t$  be greater than  $1 + \min\{v_5(c), v_2(c)\}$ , and call  $j$  the last nonzero digit of  $c$  (e.g., if  $c = 940030$  is given, it follows that  $j(940030) = 3$ ). Then,  $V((10^{k+t} + 10^{t-v_5(c)} + 1)^c) = t$  if  $c : j \nmid 5$  (by [3], Equation (16), line 7) and  $V((10^{k+t} + 10^{t-v_2(c)} + 1)^c) = t$  otherwise (by [3], Equation (16), line 8). In particular, if  $2 \nmid c$  or  $5 \nmid c$ , then  $V((10^{k+c} + 10^c + 1)^c) = c$  (since  $\min\{v_5(c), v_2(c)\} = 0 \Leftrightarrow 2 \nmid c \vee 5 \nmid c$ ).*

#### 4. Conclusion

In the previous section, for each  $c \in \mathbb{N}$ , we have provided the general equation  $V((10^{k+t} + 10^{t-\min\{v_5(c), v_2(c)\}} + 1)^c) = t$  ( $k, t \in \mathbb{N} : t > \min\{v_5(c), v_2(c)\} + 1$ ) which shows the existence of infinitely many perfect powers of degree  $c$  with a constant congruence speed of  $\min\{v_5(c), v_2(c)\} + 2$ ,  $\min\{v_5(c), v_2(c)\} + 3$ ,  $\min\{v_5(c), v_2(c)\} + 4$ ,  $\min\{v_5(c), v_2(c)\} + 5$ , and so forth.

In conclusion, for any given positive integer  $c$ , we have constructed an infinite set of  $c$ -th perfect powers that are also characterized by a constant congruence speed of  $c$ .



**Acknowledgments:** The author would like to thank his friend Aldo Roberto Pessolano for having shared with him his first thoughts about the outcome of the computer search performed on the smallest integers of the form  $a^c$ , congruent to 5 modulo 20, and characterized by a constant congruence speed of  $c$ .

**Conflict of interest:** The author declares no conflict of interest.

## References

1. Googology Wiki, Tetration. Available online: <https://googology.wikia.org/wiki/Tetration> (accessed on 31 October 2024).
2. Ripà M. The congruence speed formula. Notes Number Theory Discrete Math. 2021; 27(4): 43–61. doi:10.7546/nntdm.2021.27.4.43-61.
3. Ripà M, Onnis L. Number of stable digits of any integer tetration, Notes Number Theory Discrete Math. 2022; 28(3): 441–457. doi:10.7546/nntdm.2022.28.3.441-457.
4. The On-Line Encyclopedia of Integer Sequences (OEIS). Available online: <https://oeis.org/> (accessed on 31 October 2024).
5. Automorphic Number [From MathWorld A Wolfram Web Resource by Weisstein EW]. Available online: <https://mathworld.wolfram.com/AutomorphicNumber.html> (accessed on 31 October 2024).
6. deGuerre V, Fairbairn RA. Automorphic numbers. J. Rec. Math. 1968; 1(3): 173–179.
7. Lewis RY. A formal proof of Hensel’s lemma over the  $p$ -adic integers. In: Proceedings of the 8th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2019); 14–15 January 2019; Cascais, Portugal.
8. Germain J. On the Equation  $a^x \equiv x \pmod{b}$ . Integers. 2009; 9(6): 629–638. doi:10.1515/INTEG.2009.050.
9. Urroz JJ, Yebra JLA. On the Equation  $a^x \equiv x \pmod{b^n}$ . J. Integer Seq. 2009; 12(8): Article 09.8.8.
10. Quick L.  $p$ -adic Absolute Values [2020 Summer Program for Undergraduates]. REU 2020-The University of Chicago Mathematics; 2020.
11. Parvardi AH. Lifting The Exponent Lemma (LTE)-Version 6. Available online: <http://s3.amazonaws.com/aops-cdn.artofproblemsolving.com/resources/articles/lifting-the-exponent.pdf> (accessed on 15 May 2024).
12. Heuberger C, Mazzoli M. Elliptic curves with isomorphic groups of points over finite field extensions. J. Number Theory. 2017; 181: 89–98. doi: 10.1016/j.jnt.2017.05.028.
13. Cassels JWS. Local Fields. London Mathematical Society Student Texts 3. Cambridge University Press; 1986.