

Positivity results to iterative system of higher order boundary value problems

Kanakayya Namana¹, Sreedhar Namburi^{1,*}, Rajendra Prasad Kapula²

¹ Department of Mathematics, School of Science, GITAM (Deemed to be University), Visakhapatnam 530045, India

² Department of Applied Mathematics, College of Science and Technology, Andhra University, Visakhapatnam 530003, India

* Corresponding author: Sreedhar Namburi, sreedharnamburi@rediffmail.com

CITATION

Namana K, Namburi S, Kapula RP. Positivity results to iterative system of higher order boundary value problems. *Journal of AppliedMath*. 2024; 2(5): 1829. <https://doi.org/10.59400/jam1829>

ARTICLE INFO

Received: 8 October 2024

Accepted: 23 October 2024

Available online: 30 October 2024

COPYRIGHT



Copyright © 2024 author(s).

Journal of AppliedMath is published by Academic Publishing Pte. Ltd.

This work is licensed under the Creative Commons Attribution (CC BY) license.

<https://creativecommons.org/licenses/by/4.0/>

Abstract: The present research explores the existence of positive solutions for the iterative system of higher-order differential equations with integral boundary conditions that include a non-homogeneous term. To address the boundary value problem, the solution is expressed as a solution of an equivalent integral equation involving kernels. Subsequently, bounds for these kernels are determined to facilitate further analysis. The primary tool employed in this study is the Guo-Krasnosel'skii fixed-point theorem, which is utilized to establish the existence of positive solutions within a cone of a Banach space. This approach enables a rigorous exploration of the existence of at least one positive solution and provides insights into the behavior of the differential equation under the given boundary conditions.

Keywords: iterative system; non-homogenous conditions; positivity results; kernel

1. Introduction

Differential equations are useful for formulating mathematical models to analyze real-world phenomena. These models are frequently expressed as either initial or boundary value problems. The problems with integral boundary conditions are commonly encountered in various fields of science and engineering, such as thermoelectricity, thermoelasticity, plasma physics, underground water flow, hydrodynamic problems, chemical engineering, and many more; see Cannon [1], Ionkin [2], Chegis [3]. We refer to the study on differential equations of third order, fourth order, and higher order with integral boundary conditions [4–10].

Complex structures with multiple degrees of freedom are frequently described by systems of differential equations under specific conditions. The main challenge is to analyze mathematical models for such structures and to determine the positive solutions using various mathematical techniques. Due to the theoretical and practical significance of this topic, researchers have shown considerable interest in studying positive solutions for iterative systems of nonlinear boundary value problems by determining the parameter intervals. A few papers along these lines include Henderson et al. [11], Henderson et al. [12], Prasad et al. [13], and Oz and Karaca [14] for second-order systems; Bouteraa et al. [15] for fourth-order systems; and Henderson and Ntouyas [16], Prasad et al. [17], and Namburi et al. [18] for n th-order systems. In particular, iterative differential equations provide a novel approach to studying functional differential equations [19,20]. However, only a few studies in the literature address the existence of positive solutions for iterative systems of nonlinear boundary value problems that do not involve parameters [21,22]. In this work, we extend these

results to iterative systems of higher-order boundary value problems involving sets of equations.

Based on the above literature, we investigate the presence of positive solutions to the following iterative system of nonlinear boundary value problems,

$$\left. \begin{aligned} w_i^{(\beta)}(\varpi) + a_i(\varpi)f_i(w_{i+1}(\varpi)) &= 0, \quad 1 \leq i \leq m, \varpi \in [0,1], \\ w_{m+1}(\varpi) &= w_1(\varpi), \varpi \in [0,1], \end{aligned} \right\} \quad (1)$$

fulfilling conditions with non-homogeneous term

$$\left. \begin{aligned} w_i(0) = 0, w'_i(0) = 0, \dots, w_i^{(\beta-2)}(0) &= 0, \\ w_i^{(\gamma)}(1) - \eta_i \int_0^1 g_i(\xi) w_i^{(\gamma)}(\xi) d\xi &= \beta_i, \quad 1 \leq i \leq m, \end{aligned} \right\} \quad (2)$$

where γ is a fixed value ranging from 1 to $\beta - 2$, $\beta \geq 3$, η_i is a positive real constant, $\beta_i \in (0, \infty)$ is a parameter for $1 \leq i \leq m$, by fixed point theorem of Guo-Krasnosel'skii. By giving different values to the constants β , η_i and β_i , we get various order problems, for instance [23,24]. Here, the integral kernel plays a key role in establishing our results. The applications of integral equations are vast, spanning areas such as signal processing, fluid flow, electromagnetics, quantum mechanics, and population dynamics. However, solving integral equations presents challenges due to kernel singularity, high computational costs for high-dimensional systems, and convergence difficulties in nonlinear cases. Advancements in numerical methods continue to expand the applicability of integral equations, providing new insights and solutions to complex real-world problems [25–29].

For $1 \leq i \leq m$, the below stated assumptions are valid in this paper:

(E1) $g_i \in C([0,1], \mathbb{R}^+)$, $f_i \in C(\mathbb{R}^+, \mathbb{R}^+)$, $a_i(\varpi) \in C([0,1], \mathbb{R}^+)$ and $a_i(\varpi)$ fails to vanish identically on any closed subset of $[0,1]$,

(E2) $\forall_i = (\beta - 1)! (1 - \eta_i \theta_i) > 0$, where $\theta_i = \int_0^1 g_i(\zeta) \zeta^{\beta-\gamma-1} d\zeta$.

Define the nonnegative extended real numbers f_{i0} and $f_{i\infty}$ as $f_{i0} = \lim_{w \rightarrow 0^+} \frac{f_i(w)}{w}$

and $f_{i\infty} = \lim_{w \rightarrow \infty} \frac{f_i(w)}{w}$, for $1 \leq i \leq m$.

1.1. Definition

By a positive solution of the problem (1) and (2), we mean that $(w_1(\varpi), w_2(\varpi), \dots, w_m(\varpi)) \in (C^\beta[0,1])^m$ satisfying the Equations (1) and (2) with $w_i(\varpi) \geq 0$, $i = 1, 2, \dots, m$ for $\varpi \in [0,1]$ and $(w_1(\varpi), w_2(\varpi), \dots, w_m(\varpi)) \neq (0, 0, \dots, 0)$.

The remaining part of the paper is structured as follows: In section 2, the solution for Equations (1) and (2) is represented as a solution of the related integral equation, which includes kernels. Section 3 is devoted to studying the presence of positive solutions to Equations (1) and (2) based on the Guo-Krasnosel'skii theorem. The established results are verified by constructing the examples in section 4. The last section presents the conclusion and future scope of the study.

2. Inequalities for kernels

In this section, we represent the solution of the Equations (1) and (2) as a solution of the related integral equation with kernels. We then derive some inequalities related to these kernels.

2.1. Lemma [18]

Suppose that (E2) is true. Let $Q(x) \in C([0,1], \mathbb{R}^+)$. Then the solution of the differential equation

$$w_i^{(\beta)}(x) + Q(x) = 0, \quad 1 \leq i \leq m, x \in [0,1], \tag{3}$$

fulfilling the Equation (2) is expressed uniquely as

$$w_i(x) = \frac{\beta_i(\beta - \gamma - 1)! x^{\beta-1}}{\Upsilon_i} + \int_0^1 \left[R(x, \nu) + \frac{\eta_i x^{\beta-1}}{\Upsilon_i} \int_0^1 \zeta(\xi, \nu) g_i(\xi) d\xi \right] Q(\nu) d\nu, \tag{4}$$

$$R(x, \nu) = \frac{1}{(\beta - 1)!} \begin{cases} [x^{\beta-1}(1 - \nu)^{\beta-\gamma-1} - (x - \nu)^{\beta-1}], & 0 \leq \nu \leq x \leq 1, \\ x^{\beta-1}(1 - \nu)^{\beta-\gamma-1}, & 0 \leq x \leq \nu \leq 1, \end{cases} \tag{5}$$

and

$$\zeta(\xi, \nu) = \begin{cases} \xi^{\beta-\gamma-1}(1 - \nu)^{\beta-\gamma-1} - (\xi - \nu)^{\beta-\gamma-1}, & 0 \leq \nu \leq \xi \leq 1, \\ \xi^{\beta-\gamma-1}(1 - \nu)^{\beta-\gamma-1}, & 0 \leq \xi \leq \nu \leq 1. \end{cases} \tag{6}$$

2.2. Lemma [18]

The functions $R(x, \nu)$ and $\zeta(x, \nu)$ satisfy the below:

- (1) $R(x, \nu)$ and $\zeta(x, \nu)$ are nonnegative, for every $x, \nu \in [0,1]$,
- (2) $R(x, \nu) \leq R(1, \nu)$, for every $x, \nu \in [0,1]$,
- (3) $\frac{1}{4^{\beta-1}} R(1, \nu) \leq R(x, \nu)$, for every $x \in I$ and $\nu \in [0,1]$, where $I = [\frac{1}{4}, \frac{3}{4}]$.

We can see that the solution for the Equations (1) and (2) is an m -tuple $(w_1(x), w_2(x), \dots, w_m(x))$ if and only if $w_i(x)$ fulfills the subsequent equations

$$w_i(x) = \frac{\beta_i(\beta - \gamma - 1)! x^{\beta-1}}{\Upsilon_i} + \int_0^1 \left[R(x, \nu) + \frac{\eta_i x^{\beta-1}}{\Upsilon_i} \int_0^1 \zeta(\xi, \nu) g_i(\xi) d\xi \right] a_i(\nu) f_i(w_{i+1}(\nu)) d\nu, \tag{7}$$

$$1 \leq i \leq m, x \in [0,1],$$

and

$$w_{m+1}(x) = w_1(x), x \in [0,1].$$

Therefore,

$$w_1(x) = \frac{\beta_1(\beta - \gamma - 1)! x^{\beta-1}}{\Upsilon_1} + \int_0^1 \left[R(x, \nu_1) + \frac{\eta_1 x^{\beta-1}}{\Upsilon_1} \int_0^1 \zeta(\xi, \nu_1) g_1(\xi) d\xi \right] a_1(\nu_1) f_1 \left(\frac{\beta_2(\beta - \gamma - 1)! \nu_1^{\beta-1}}{\Upsilon_2} + \int_0^1 \left[R(\nu_1, \nu_2) + \frac{\eta_2 \nu_1^{\beta-1}}{\Upsilon_2} \int_0^1 \zeta(\xi, \nu_2) g_2(\xi) d\xi \right] \right) d\nu_1$$

$$\begin{aligned}
 & a_2(v_2) \dots f_{m-1} \left(\frac{\beta_m (\beta - r - 1)! v_{m-1}^{\beta-1}}{\gamma_m} \right. \\
 & + \int_0^1 \left[R(v_{m-1}, v_m) \right. \\
 & \left. \left. + \frac{\eta_m v_{m-1}^{\beta-1}}{\gamma_m} \int_0^1 \zeta(\xi, v_m) g_m(\xi) d\xi \right] a_m(v_m) f_m(w_1(v_m)) dv_m \right) \dots dv_2 \Big) dv_1.
 \end{aligned}$$

To establish main outcomes, we shall use the Guo-Krasnosel'skii fixed point theorem, which is described below.

2.3. Theorem [30,31]

Let ρ be a cone in a Banach Space \mathbb{B} . Let the two open subsets be Λ_1 and Λ_2 of Banach space \mathbb{B} such that $0 \in \Lambda_1$ and $\overline{\Lambda_1} \subset \Lambda_2$. If the function $T: \rho \cap (\overline{\Lambda_2} \setminus \Lambda_1) \rightarrow \rho$ satisfy the below subsequent inequalities:

- (i) $\|Tu\| \leq \|u\|$, for $u \in \rho \cap \partial\Lambda_1$ and $\|Tu\| \geq \|u\|$, for $u \in \rho \cap \partial\Lambda_2$, or
- (ii) $\|Tu\| \geq \|u\|$, for $u \in \rho \cap \partial\Lambda_1$ and $\|Tu\| \leq \|u\|$, for $u \in \rho \cap \partial\Lambda_2$,

then there is a fixed point in $\rho \cap (\overline{\Lambda_2} \setminus \Lambda_1)$.

3. Positive solutions

This section contains the presence of positive solutions to the Equations (1) and (2). For construction, let the set $\mathbb{B} = \{w : w \in C([0,1], \mathbb{R})\}$ be a Banach space with

$$\|w\| = \max_{x \in [0,1]} |w(x)|.$$

Let ρ be a cone in a Banach space \mathbb{B} and is defined as

$$\rho = \{w \in \mathbb{B} : w(x) \geq 0, \text{ for every } x \in [0,1] \text{ with } \min_{x \in I} w(x) \geq \frac{1}{4^{\beta-1}} \|w\|\}.$$

For $w_1 \in \rho$, define a function $\mathfrak{D}: \rho \rightarrow \mathbb{B}$ as

$$\mathfrak{D}w_1(x) = \left. \begin{aligned}
 & \frac{\beta_1 (\beta - r - 1)! x^{\beta-1}}{\gamma_1} + \int_0^1 \left[R(x, v_1) + \frac{\eta_1 x^{\beta-1}}{\gamma_1} \int_0^1 \zeta(\xi, v_1) g_1(\xi) d\xi \right] \\
 & a_1(v_1) f_1 \left(\frac{\beta_2 (\beta - r - 1)! v_1^{\beta-1}}{\gamma_2} + \int_0^1 \left[R(v_1, v_2) + \frac{\eta_2 v_1^{\beta-1}}{\gamma_2} \int_0^1 \zeta(\xi, v_2) g_2(\xi) d\xi \right] \right. \\
 & a_2(v_2) \dots f_{m-1} \left(\frac{\beta_m (\beta - r - 1)! v_{m-1}^{\beta-1}}{\gamma_m} + \int_0^1 \left[R(v_{m-1}, v_m) + \frac{\eta_m v_{m-1}^{\beta-1}}{\gamma_m} \int_0^1 \zeta(\xi, v_m) g_m(\xi) d\xi \right] \right. \\
 & \left. \left. \left. \left. a_m(v_m) f_m(w_1(v_m)) dv_m \right) \dots dv_2 \right) dv_1. \right. \right. \right. \right.
 \end{aligned} \right\} \tag{7}$$

3.1. Lemma

The function $\mathfrak{D}: \rho \rightarrow \mathfrak{B}$ stated in Equation (7) is a self-map on ρ .

Proof. For $w_1 \in \rho$, $\mathfrak{D}w_1(x) \geq 0$ on $x \in [0,1]$ by the nonnegativity of $R(x, v)$ and $\mathfrak{S}(x, v)$. Using Lemma 2.2, we see that for $w_1 \in \rho$,

$$\begin{aligned} \mathfrak{D}w_1(x) &= \frac{\beta_1(\beta-\gamma-1)!x^{\beta-1}}{\Upsilon_1} + \int_0^1 \left[R(x, v_1) + \frac{\eta_1 x^{\beta-1}}{\Upsilon_1} \int_0^1 \mathfrak{S}(\xi, v_1)g_1(\xi)d\xi \right] \\ a_1(v_1)f_1 &\left(\frac{\beta_2(\beta-\gamma-1)!v_1^{\beta-1}}{\Upsilon_2} + \int_0^1 \left[R(v_1, v_2) + \frac{\eta_2 v_1^{\beta-1}}{\Upsilon_2} \int_0^1 \mathfrak{S}(\xi, v_2)g_2(\xi)d\xi \right] \right. \\ a_2(v_2) \dots f_{m-1} &\left(\frac{\beta_m(\beta-\gamma-1)!v_{m-1}^{\beta-1}}{\Upsilon_m} \right. \\ &\quad \left. + \int_0^1 \left[R(v_{m-1}, v_m) + \frac{\eta_m v_{m-1}^{\beta-1}}{\Upsilon_m} \int_0^1 \mathfrak{S}(\xi, v_m)g_m(\xi)d\xi \right] \right. \\ &\quad \left. a_m(v_m)f_m(w_1(v_m)) dv_m \right) \dots dv_2 \Big) dv_1 \\ &\leq \frac{\beta_1(\beta-\gamma-1)!}{\Upsilon_1} + \int_0^1 \left[R(1, v_1) + \frac{\eta_1}{\Upsilon_1} \int_0^1 \mathfrak{S}(\xi, v_1)g_1(\xi)d\xi \right] \\ a_1(v_1)f_1 &\left(\frac{\beta_2(\beta-\gamma-1)!v_1^{\beta-1}}{\Upsilon_2} + \int_0^1 \left[R(v_1, v_2) + \frac{\eta_2 v_1^{\beta-1}}{\Upsilon_2} \int_0^1 \mathfrak{S}(\xi, v_2)g_2(\xi)d\xi \right] \right. \\ a_2(v_2) \dots f_{m-1} &\left(\frac{\beta_m(\beta-\gamma-1)!v_{m-1}^{\beta-1}}{\Upsilon_m} \right. \\ &\quad \left. + \int_0^1 \left[R(v_{m-1}, v_m) + \frac{\eta_m v_{m-1}^{\beta-1}}{\Upsilon_m} \int_0^1 \mathfrak{S}(\xi, v_m)g_m(\xi)d\xi \right] \right. \\ &\quad \left. a_m(v_m)f_m(w_1(v_m)) dv_m \right) \dots dv_2 \Big) dv_1. \end{aligned}$$

Then

$$\| \mathfrak{D}w_1(x) \| \leq \left. \begin{aligned} &\frac{\beta_1(\beta-\gamma-1)!}{\Upsilon_1} + \int_0^1 \left[R(1, v_1) + \frac{\eta_1}{\Upsilon_1} \int_0^1 \mathfrak{S}(\xi, v_1)g_1(\xi)d\xi \right] \\ &a_1(v_1)f_1 \left(\frac{\beta_2(\beta-\gamma-1)!v_1^{\beta-1}}{\Upsilon_2} + \right. \\ &\quad \left. \int_0^1 \left[R(v_1, v_2) + \frac{\eta_2 v_1^{\beta-1}}{\Upsilon_2} \int_0^1 \mathfrak{S}(\xi, v_2)g_2(\xi)d\xi \right] \right. \\ &\quad a_2(v_2) \dots f_{m-1} \left(\frac{\beta_m(\beta-\gamma-1)!v_{m-1}^{\beta-1}}{\Upsilon_m} + \right. \\ &\quad \left. \int_0^1 \left[R(v_{m-1}, v_m) + \frac{\eta_m v_{m-1}^{\beta-1}}{\Upsilon_m} \int_0^1 \mathfrak{S}(\xi, v_m)g_m(\xi)d\xi \right] \right. \\ &\quad \left. a_m(v_m)f_m(w_1(v_m)) dv_m \right) \dots dv_2 \Big) dv_1. \end{aligned} \right\} \tag{8}$$

Next, if $w_1 \in \rho$, we have from Lemma 2.2 and Equation (8) that

$$\min_{x \in I} \mathfrak{D}w_1(x) = \min_{x \in I} \left\{ \frac{\beta_1(\beta-\gamma-1)!x^{\beta-1}}{\Upsilon_1} + \int_0^1 \left[R(x, v_1) + \frac{\eta_1 x^{\beta-1}}{\Upsilon_1} \int_0^1 \mathfrak{S}(\xi, v_1)g_1(\xi)d\xi \right] \right\}$$

$$\begin{aligned}
 & a_1(v_1)f_1\left(\frac{\beta_2(\beta-\gamma-1)!v_1^{\beta-1}}{\Upsilon_2} + \int_0^1\left[R(v_1,v_2) + \frac{\eta_2v_1^{\beta-1}}{\Upsilon_2}\int_0^1\zeta(\xi,v_2)g_2(\xi)d\xi\right]\right. \\
 & \quad \left. a_2(v_2)\dots f_{m-1}\left(\frac{\beta_m(\beta-\gamma-1)!v_{m-1}^{\beta-1}}{\Upsilon_m}\right.\right. \\
 & \quad \left. + \int_0^1\left[R(v_{m-1},v_m) + \frac{\eta_mv_{m-1}^{\beta-1}}{\Upsilon_m}\int_0^1\zeta(\xi,v_m)g_m(\xi)d\xi\right]\right. \\
 & \quad \left. a_m(v_m)f_m(w_1(v_m))dv_m\right)\dots dv_2\left.)dv_1\right\} \\
 & \geq \frac{1}{4^{\beta-1}}\left\{\frac{\beta_1(\beta-\gamma-1)!}{\Upsilon_1} + \int_0^1\left[R(1,v_1) + \frac{\eta_1}{\Upsilon_1}\int_0^1\zeta(\xi,v_1)g_1(\xi)d\xi\right]\right. \\
 & a_1(v_1)f_1\left(\frac{\beta_2(\beta-\gamma-1)!v_1^{\beta-1}}{\Upsilon_2} + \int_0^1\left[R(v_1,v_2) + \frac{\eta_2v_1^{\beta-1}}{\Upsilon_2}\int_0^1\zeta(\xi,v_2)g_2(\xi)d\xi\right]\right. \\
 & \quad \left. a_2(v_2)\dots f_{m-1}\left(\frac{\beta_m(\beta-\gamma-1)!v_{m-1}^{\beta-1}}{\Upsilon_m}\right.\right. \\
 & \quad \left. + \int_0^1\left[R(v_{m-1},v_m) + \frac{\eta_mv_{m-1}^{\beta-1}}{\Upsilon_m}\int_0^1\zeta(\xi,v_m)g_m(\xi)d\xi\right]\right. \\
 & \quad \left. a_m(v_m)f_m(w_1(v_m))dv_m\right)\dots dv_2\left.)dv_1\right\} \\
 & \geq \frac{1}{4^{\beta-1}}\|\mathfrak{D}w_1(\infty)\|.
 \end{aligned}$$

Hence, $\mathfrak{D} : \rho \rightarrow \rho$. This completes the proof.

In addition, we can see that the function \mathfrak{D} is completely continuous by applying the Arzela-Ascoli theorem [32].

3.2. Theorem

Suppose that (E1) and (E2) are met. If $f_{i0} = 0$ and $f_{i\infty} = \infty$ are true, then the Equations (1) and (2) has at least one positive solution and $\beta_i \in (0, \infty)$ small enough for $1 \leq i \leq m$.

Proof. For $1 \leq i \leq m$, by using the definition of $f_{i0} = 0$, there exist $\varrho_i > 0$ and $H_1 > 0$ such that

$$f_i(w) \leq \varrho_i w, \text{ for } 0 < w \leq H_1,$$

where ϱ_i satisfies

$$\varrho_i \int_0^1 2 \left[R(1, v_i) + \frac{\eta_i}{\Upsilon_i} \int_0^1 \zeta(\xi, v_i) g_i(\xi) d\xi \right] a_i(v_i) dv_i \leq 1. \tag{9}$$

For $1 \leq i \leq m$, let β_i be chosen

$$0 < \beta_i \leq \frac{\Upsilon_i H_1}{(\beta-\gamma-1)! 2}.$$

Let $w_1 \in \rho$ with $\|w_1\| = H_1$. Then using Lemma 2.2 and for $0 \leq v_{m-1} \leq 1$, we have

$$\begin{aligned} & \frac{\beta_m(\beta - \varkappa - 1)! \nu_{m-1}^{\beta-1}}{\Upsilon_m} \\ & + \int_0^1 \left[R(\nu_{m-1}, \nu_m) + \frac{\eta_m \nu_{m-1}^{\beta-1}}{\Upsilon_m} \int_0^1 \zeta(\xi, \nu_m) g_m(\xi) d\xi \right] a_m(\nu_m) f_m(w_1(\nu_m)) d\nu_m \\ & \leq \frac{\beta_m(\beta - \varkappa - 1)!}{\Upsilon_m} \\ & \quad + \int_0^1 \left[R(1, \nu_m) + \frac{\eta_m}{\Upsilon_m} \int_0^1 \zeta(\xi, \nu_m) g_m(\xi) d\xi \right] a_m(\nu_m) \varrho_m w_1(\nu_m) d\nu_m \\ & \leq \frac{H_1}{2} + \varrho_m \int_0^1 \left[R(1, \nu_m) + \frac{\eta_m}{\Upsilon_m} \int_0^1 \zeta(\xi, \nu_m) g_m(\xi) d\xi \right] a_m(\nu_m) d\nu_m \|w_1\| \\ & \leq \frac{H_1}{2} + \frac{H_1}{2} = H_1. \end{aligned}$$

It comes in the same way as Lemma 2.2 and for $0 \leq \vartheta_{m-2} \leq 1$,

$$\begin{aligned} & \frac{\beta_{m-1}(\beta - \varkappa - 1)! \nu_{m-2}^{\beta-1}}{\Upsilon_{m-1}} + \int_0^1 \left[R(\nu_{m-2}, \nu_{m-1}) + \frac{\eta_{m-1} \nu_{m-2}^{\beta-1}}{\Upsilon_{m-1}} \int_0^1 \zeta(\xi, \nu_{m-1}) g_{m-1}(\xi) d\xi \right] \\ & a_{m-1}(\nu_{m-1}) f_{m-1} \left(\frac{\beta_m(\beta - \varkappa - 1)! \nu_{m-1}^{\beta-1}}{\Upsilon_m} + \int_0^1 \left[R(\nu_{m-1}, \nu_m) + \frac{\eta_m \nu_{m-1}^{\beta-1}}{\Upsilon_m} \int_0^1 \zeta(\xi, \nu_m) g_m(\xi) d\xi \right] \right. \\ & \quad \left. a_m(\nu_m) f_m(w_1(\nu_m)) d\nu_m \right) d\nu_{m-1} \\ & \leq \frac{\beta_{m-1}(\beta - \varkappa - 1)!}{\Upsilon_{m-1}} + \int_0^1 \left[R(1, \nu_{m-1}) + \frac{\eta_{m-1}}{\Upsilon_{m-1}} \int_0^1 \zeta(\xi, \nu_{m-1}) g_{m-1}(\xi) d\xi \right] a_{m-1}(\nu_{m-1}) d\nu_{m-1} \varrho_{m-1} H_1 \\ & \leq \frac{H_1}{2} + \frac{H_1}{2} = H_1. \end{aligned}$$

Applying the similar argument, one can obtain, for $0 \leq \varkappa \leq 1$,

$$\begin{aligned} & \frac{\beta_1(\beta - \varkappa - 1)! \varkappa^{\beta-1}}{\Upsilon_1} + \int_0^1 \left[R(\varkappa, \nu_1) + \frac{\eta_1 \varkappa^{\beta-1}}{\Upsilon_1} \int_0^1 \zeta(\xi, \nu_1) g_1(\xi) d\xi \right] \\ & a_1(\nu_1) f_1 \left(\frac{\beta_2(\beta - \varkappa - 1)! \nu_1^{\beta-1}}{\Upsilon_2} + \int_0^1 \left[R(\nu_1, \nu_2) + \frac{\eta_2 \nu_1^{\beta-1}}{\Upsilon_2} \int_0^1 \zeta(\xi, \nu_2) g_2(\xi) d\xi \right] \right. \\ & a_2(\nu_2) \dots f_{m-1} \left(\frac{\beta_m(\beta - \varkappa - 1)! \nu_{m-1}^{\beta-1}}{\Upsilon_m} + \int_0^1 \left[R(\nu_{m-1}, \nu_m) + \frac{\eta_m \nu_{m-1}^{\beta-1}}{\Upsilon_m} \int_0^1 \zeta(\xi, \nu_m) g_m(\xi) d\xi \right] \right. \\ & \left. a_m(\nu_m) f_m(w_1(\nu_m)) d\nu_m \right) \dots d\nu_2 \left. \right) d\nu_1 \leq H_1, \end{aligned}$$

so that for $0 \leq \varkappa \leq 1$,

$$\mathfrak{D}w_1(\varkappa) \leq H_1.$$

Hence, $\|\mathfrak{D}w_1\| \leq H_1 = \|w_1\|$. Take

$$\Lambda_1 = \{w \in \mathfrak{B} : \|w\| < H_1\}.$$

Then

$$\|\mathfrak{D}w_1\| \leq \|w_1\|, \quad \text{for } w_1 \in \rho \cap \partial\Lambda_1. \tag{10}$$

Since $f_{i\infty} = \infty$, $1 \leq i \leq m$, there exist $\sigma_i > 0$ and $\overline{H}_2 \geq 0$, such that

$$f_i(w) \geq \sigma_i w, \quad \text{for } w \geq \overline{H}_2,$$

where σ_i satisfies

$$\frac{\sigma_i}{4^{2\beta-2}} \int_{\nu_i \in I} \left[R(1, \nu_i) + \frac{\eta_i}{\Upsilon_i} \int_{\xi \in I} \xi(\zeta, \nu_i) g_i(\xi) d\zeta \right] a_i(\nu_i) d\nu_i \geq 1. \tag{11}$$

Let

$$H_2 = \max \{2H_1, 4^{\beta-1}\overline{H}_2\}.$$

Choose $w_1 \in \rho$ and $\|w_1\| = H_2$. Then

$$\min_{\nu \in I} w_1(\nu) \geq \frac{1}{4^{\beta-1}} \|w_1\| \geq \overline{H}_2.$$

By Lemma 2.2 and for $\frac{1}{4} \leq \nu_{m-1} \leq \frac{3}{4}$, we have

$$\begin{aligned} & \frac{\beta_m (\beta - \nu - 1)! \nu_{m-1}^{\beta-1}}{\Upsilon_m} \\ & + \int_0^1 \left[R(\nu_{m-1}, \nu_m) + \frac{\eta_m \nu_{m-1}^{\beta-1}}{\Upsilon_m} \int_0^1 \xi(\xi, \nu_m) g_m(\xi) d\xi \right] a_m(\nu_m) f_m(w_1(\nu_m)) d\nu_m \\ & \geq \int_0^1 \left[R(\nu_{m-1}, \nu_m) + \frac{\eta_m \nu_{m-1}^{\beta-1}}{\Upsilon_m} \int_0^1 \xi(\xi, \nu_m) g_m(\xi) d\xi \right] a_m(\nu_m) f_m(w_1(\nu_m)) d\nu_m \\ & \geq \frac{1}{4^{\beta-1}} \int_{\nu_m \in I} \left[R(1, \nu_m) + \frac{\eta_m}{\Upsilon_m} \int_{\xi \in I} \xi(\xi, \nu_m) g_m(\xi) d\xi \right] a_m(\nu_m) \sigma_m w_1(\nu_m) d\nu_m \\ & \geq \frac{1}{4^{2\beta-2}} \int_{\nu_m \in I} \left[R(1, \nu_m) + \frac{\eta_m}{\Upsilon_m} \int_{\xi \in I} \xi(\xi, \nu_m) g_m(\xi) d\xi \right] a_m(\nu_m) d\nu_m \sigma_m \|w_1\| \\ & \geq \|w_1\| = H_2. \end{aligned}$$

It comes in the same way as Lemma 2.2 and for $\frac{1}{4} \leq \nu_{m-1} \leq \frac{3}{4}$,

$$\begin{aligned} & \frac{\beta_{m-1} (\beta - \nu - 1)! \nu_{m-2}^{\beta-1}}{\Upsilon_{m-1}} + \int_0^1 \left[R(\nu_{m-2}, \nu_{m-1}) + \frac{\eta_{m-1} \nu_{m-2}^{\beta-1}}{\Upsilon_{m-1}} \int_0^1 \xi(\xi, \nu_{m-1}) g_{m-1}(\xi) d\xi \right] \\ & \quad a_{m-1}(\nu_{m-1}) f_{m-1} \left(\frac{\beta_m (\beta - \nu - 1)! \nu_{m-1}^{\beta-1}}{\Upsilon_m} + \int_0^1 \left[R(\nu_{m-1}, \nu_m) + \right. \right. \\ & \quad \left. \left. \frac{\eta_m \nu_{m-1}^{\beta-1}}{\Upsilon_m} \int_0^1 \xi(\xi, \nu_m) g_m(\xi) d\xi \right] a_m(\nu_m) f_m(w_1(\nu_m)) d\nu_m \right) d\nu_{m-1} \\ & \geq \frac{1}{4^{\beta-1}} \int_{\nu_{m-1} \in I} \left[R(1, \nu_{m-1}) \right. \\ & \quad \left. + \frac{\eta_{m-1}}{\Upsilon_{m-1}} \int_{\xi \in I} \xi(\xi, \nu_{m-1}) g_{m-1}(\xi) d\xi \right] a_{m-1}(\nu_{m-1}) d\nu_{m-1} \sigma_{m-1} H_2 \end{aligned}$$

$$\begin{aligned} &\geq \frac{1}{4^{2\beta-2}} \int_{\nu_{m-1} \in I} \left[R(1, \nu_{m-1}) \right. \\ &\quad \left. + \frac{\eta_{m-1}}{\gamma_{m-1}} \int_{\xi \in I} \zeta(\xi, \nu_{m-1}) g_{m-1}(\xi) d\xi \right] a_{m-1}(\nu_{m-1}) d\nu_{m-1} \sigma_{m-1} H_2 \\ &\geq H_2. \end{aligned}$$

Proceeding in a similar argument, we have

$$\begin{aligned} &\frac{\beta_1(\beta-\gamma-1)! \alpha^{\beta-1}}{\gamma_1} + \int_0^1 \left[R(\alpha, \nu_1) + \frac{\eta_1 \alpha^{\beta-1}}{\gamma_1} \int_0^1 \zeta(\xi, \nu_1) g_1(\xi) d\xi \right] \\ &a_1(\nu_1) f_1 \left(\frac{\beta_2(\beta-\gamma-1)! \nu_1^{\beta-1}}{\gamma_2} + \int_0^1 \left[R(\nu_1, \nu_2) + \frac{\eta_2 \nu_1^{\beta-1}}{\gamma_2} \int_0^1 \zeta(\xi, \nu_2) g_2(\xi) d\xi \right] \right. \\ &a_2(\nu_2) \dots f_{m-1} \left(\frac{\beta_m(\beta-\gamma-1)! \nu_{m-1}^{\beta-1}}{\gamma_m} \right. \\ &\quad \left. + \int_0^1 \left[R(\nu_{m-1}, \nu_m) + \frac{\eta_m \nu_{m-1}^{\beta-1}}{\gamma_m} \int_0^1 \zeta(\xi, \nu_m) g_m(\xi) d\xi \right] \right. \\ &\quad \left. a_m(\nu_m) f_m(w_1(\nu_m)) d\nu_m \right) \dots d\nu_2 \Big) d\nu_1 \geq H_2, \end{aligned}$$

so that, for $0 \leq \alpha \leq 1$,

$$\mathfrak{D}w_1(\alpha) \geq H_2 = \|w_1\|.$$

Hence, $\|\mathfrak{D}w_1\| \geq \|w_1\|$.

Take $\Lambda_2 = \{w \in \mathcal{B} : \|w_1\| < H_2\}$, then

$$\|\mathfrak{D}w_1\| \geq \|w_1\|, \quad \text{for } w_1 \in \rho \cap \partial\Lambda_2. \tag{12}$$

Using Theorem 2.3 to the Equations (10) and (12), the function \mathfrak{D} has a fixed point $w_1 \in \rho \cap (\overline{\Lambda_2} \setminus \Lambda_1)$ and that point is the positive solution (w_1, w_2, \dots, w_m) of the Equations (1) and (2) by taking $w_{m+1} = w_1$. Therefore, the solution can be iteratively expressed as

$$\begin{aligned} w_i(\alpha) &= \frac{\beta_i(\beta-\gamma-1)! \alpha^{\beta-1}}{\gamma_i} + \int_0^1 \left[R(\alpha, \nu) + \right. \\ &\quad \left. \frac{\eta_i \alpha^{\beta-1}}{\gamma_i} \int_0^1 \zeta(\xi, \nu) g_i(\xi) d\xi \right] a_i(\nu) f_i(w_{i+1}(\nu)) d\nu, \\ &\quad i = m, m-1, \dots, 1. \end{aligned}$$

3.3. Theorem

Suppose that (E1) and (E2) are met. If $f_{i0} = \infty$ and $f_{i\infty} = 0$ are true, then the Equations (1) and (2) has at least one positive solution and $\beta_i \in (0, \infty)$ small enough for $1 \leq i \leq m$.

Proof. For $1 \leq i \leq m$, by the definition of f_{i0} , there exist $\varepsilon_i > 0$ and $H_3 > 0$ such that

$$f_i(w) \geq \varepsilon_i w, \quad \text{for } 0 < w \leq H_3,$$

where $\varepsilon_i \geq \sigma_i$ and σ_i is given in Equation (11).

Choose $w_1 \in \rho$ and $\|w_1\| = H_3$. Then, we have from Lemma 2.2 and for $\frac{1}{4} \leq v_{m-1} \leq \frac{3}{4}$, we have

$$\begin{aligned} & \frac{\beta_m(\beta - \gamma - 1)! v_{m-1}^{\beta-1}}{\gamma_m} \\ & + \int_0^1 \left[R(v_{m-1}, v_m) \right. \\ & \left. + \frac{\eta_m v_{m-1}^{\beta-1}}{\gamma_m} \int_0^1 \zeta(\xi, v_m) g_m(\xi) d\xi \right] a_m(v_m) f_m(w_1(v_m)) dv_m \\ \geq & \int_0^1 \left[R(v_{m-1}, v_m) + \frac{\eta_m v_{m-1}^{\beta-1}}{\gamma_m} \int_0^1 \zeta(\xi, v_m) g_m(\xi) d\xi \right] a_m(v_m) f_m(w_1(v_m)) dv_m \\ \geq & \frac{1}{4^{\beta-1}} \int_{v_m \in I} \left[R(1, v_m) + \frac{\eta_m}{\gamma_m} \int_{\xi \in I} \zeta(\xi, v_m) g_m(\xi) d\xi \right] a_m(v_m) \varepsilon_m w_1(v_m) dv_m \\ \geq & \frac{1}{4^{2\beta-2}} \int_{v_m \in I} \left[R(1, v_m) + \frac{\eta_m}{\gamma_m} \int_{\xi \in I} \zeta(\xi, v_m) g_m(\xi) d\xi \right] a_m(v_m) dv_m \varepsilon_m \|w_1\| \\ & \geq \|w_1\| = H_3. \end{aligned}$$

It comes in the same way as Lemma 2.2 and for $\frac{1}{4} \leq v_{m-1} \leq \frac{3}{4}$,

$$\begin{aligned} & \frac{\beta_{m-1}(\beta - \gamma - 1)! v_{m-2}^{\beta-1}}{\gamma_{m-1}} + \int_0^1 \left[R(v_{m-2}, v_{m-1}) + \frac{\eta_{m-1} v_{m-2}^{\beta-1}}{\gamma_{m-1}} \int_0^1 \zeta(\xi, v_{m-1}) g_{m-1}(\xi) d\xi \right] \\ & a_{m-1}(v_{m-1}) f_{m-1} \left(\frac{\beta_m(\beta - \gamma - 1)! v_{m-1}^{\beta-1}}{\gamma_m} + \int_0^1 \left[R(v_{m-1}, v_m) + \right. \right. \\ & \left. \left. \frac{\eta_m v_{m-1}^{\beta-1}}{\gamma_m} \int_0^1 \zeta(\xi, v_m) g_m(\xi) d\xi \right] a_m(v_m) f_m(w_1(v_m)) dv_m \right) dv_{m-1} \\ \geq & \frac{1}{4^{\beta-1}} \int_{v_{m-1} \in I} \left[R(1, v_{m-1}) \right. \\ & \left. + \frac{\eta_{m-1}}{\gamma_{m-1}} \int_{\xi \in I} \zeta(\xi, v_{m-1}) g_{m-1}(\xi) d\xi \right] a_{m-1}(v_{m-1}) dv_{m-1} \varepsilon_{m-1} H_3 \\ \geq & \frac{1}{4^{2\beta-2}} \int_{v_{m-1} \in I} \left[R(1, v_{m-1}) \right. \\ & \left. + \frac{\eta_{m-1}}{\gamma_{m-1}} \int_{\xi \in I} \zeta(\xi, v_{m-1}) g_{m-1}(\xi) d\xi \right] a_{m-1}(v_{m-1}) dv_{m-1} \varepsilon_{m-1} H_3 \\ \geq & H_3. \end{aligned}$$

Using a similar argument, we have

$$\begin{aligned} & \frac{\beta_1(\beta - \gamma - 1)! \varkappa^{\beta-1}}{\gamma_1} + \int_0^1 \left[R(\varkappa, v_1) + \frac{\eta_1 \varkappa^{\beta-1}}{\gamma_1} \int_0^1 \zeta(\xi, v_1) g_1(\xi) d\xi \right] \\ & a_1(v_1) f_1 \left(\frac{\beta_2(\beta - \gamma - 1)! v_1^{\beta-1}}{\gamma_2} + \int_0^1 \left[R(v_1, v_2) + \frac{\eta_2 v_1^{\beta-1}}{\gamma_2} \int_0^1 \zeta(\xi, v_2) g_2(\xi) d\xi \right] \right) \end{aligned}$$

$$a_2(\nu_2) \dots f_{m-1} \left(\frac{\beta_m(\beta - \gamma - 1)! \nu_{m-1}^{\beta-1}}{\gamma_m} + \int_0^1 \left[R(\nu_{m-1}, \nu_m) + \frac{\eta_m \nu_{m-1}^{\beta-1}}{\gamma_m} \int_0^1 \zeta(\xi, \nu_m) g_m(\xi) d\xi \right] a_m(\nu_m) f_m(w_1(\nu_m)) d\nu_m \right) \dots d\nu_2 \Big) d\nu_1 \geq H_3,$$

so that

$$\mathcal{D}w_1 \geq H_3 = \|w_1\|.$$

Hence, $\|\mathcal{D}w_1\| \geq \|w_1\|$. If we put

$$\Lambda_3 = \{w \in \mathcal{B} : \|w_1\| < H_3\},$$

then

$$\|\mathcal{D}w_l\| \geq \|w_l\|, \quad \text{for } w_l \in \rho \cap \partial\Lambda_3. \tag{13}$$

Since $f_{i\infty} = 0$, for $1 \leq i \leq m$, there exist $\zeta_i > 0$ and $\bar{H}_4 > 0$ such that

$$f_i(w) \leq \zeta_i w, \text{ for } w \geq \bar{H}_4,$$

where $\zeta_i \leq \varrho_i$ and ϱ_i is given in Equation (9).

For $1 \leq i \leq m$, set

$$f_i^*(w) = \sup_{0 \leq s \leq w} f_i(s).$$

It is evident from the fact that for $1 \leq i \leq m$, the real-valued function $f_i^*(w)$ is non-decreasing, $f_i \leq f_i^*$ and

$$\lim_{w \rightarrow \infty} \frac{f_i^*(w)}{w} = 0.$$

As a result, for $1 \leq i \leq m$, there exists $H_4 > \max\{2H_3, \bar{H}_4\}$ such that

$$f_i^*(w) \leq f_i^*(H_4), \quad 0 < w \leq H_4.$$

Let $\beta_i, 1 \leq i \leq m$, fulfills

$$0 < \beta_i \leq \frac{\gamma_i H_4}{(\beta - \gamma - 1)!}.$$

Choose $w_1 \in \rho$ and $\|w_1\| = H_4$. Apply same argument repeatedly, we have

$$\begin{aligned} \mathcal{D}w_1(x) &= \frac{\beta_1(\beta - \gamma - 1)! x^{\beta-1}}{\gamma_1} + \int_0^1 \left[R(x, \nu_1) + \frac{\eta_1 x^{\beta-1}}{\gamma_1} \int_0^1 \zeta(\xi, \nu_1) g_1(\xi) d\xi \right] \\ a_1(\nu_1) f_1 \left(\frac{\beta_2(\beta - \gamma - 1)! \nu_1^{\beta-1}}{\gamma_2} + \int_0^1 \left[R(\nu_1, \nu_2) + \frac{\eta_2 \nu_1^{\beta-1}}{\gamma_2} \int_0^1 \zeta(\xi, \nu_2) g_2(\xi) d\xi \right] \right. \\ &\quad \left. a_2(\nu_2) \dots f_m(w_1(\nu_m)) d\nu_m \right) \dots d\nu_2 \Big) d\nu_1 \\ &\leq \frac{\beta_1(\beta - \gamma - 1)!}{\gamma_1} + \int_0^1 \left[R(1, \nu_1) + \frac{\eta_1}{\gamma_1} \int_0^1 \zeta(\xi, \nu_1) g_1(\xi) d\xi \right] \end{aligned}$$

$$\begin{aligned}
 & a_1(v_1)f_1^* \left(\frac{\beta_2(\beta - \gamma - 1)! v_1^{\beta-1}}{\gamma_2} + \int_0^1 \left[R(v_1, v_2) + \frac{\eta_2 v_1^{\beta-1}}{\gamma_2} \int_0^1 \zeta(\xi, v_2)g_2(\xi)d\xi \right] \right. \\
 & \quad \left. a_2(v_2) \dots f_m(w_1(v_m)) dv_m \dots dv_2 \right) dv_1 \\
 & \leq \frac{H_4}{2} + \int_0^1 \left[R(1, v_1) + \frac{\eta_1}{\gamma_1} \int_0^1 \zeta(\xi, v_1)g_1(\xi) d\xi \right] a_1(v_1)f_1^*(H_4)dv \\
 & \leq \frac{H_4}{2} + \int_0^1 \left[R(1, v_1) + \frac{\eta_1}{\gamma_1} \int_0^1 \zeta(\xi, v_1)g_1(\xi) d\xi \right] a_1(v_1)dv_1 \zeta_1 H_4 \\
 & \leq \frac{H_4}{2} + \frac{H_4}{2} = H_4.
 \end{aligned}$$

Hence, $\|\mathfrak{D}w_1\| \leq \|w_1\|$. Thus, we choose

$$\Lambda_4 = \{w \in \mathbb{B} : \|w_1\| < H_4\},$$

then

$$\|\mathfrak{D}w_l\| \leq \|w_l\|, \quad \text{for } w_l \in \rho \cap \partial\Lambda_4. \tag{14}$$

Using Theorem 2.3 to the Equations (13) and (14), it can see that \mathfrak{D} has a fixed point $w_1 \in \rho \cap (\overline{\Lambda_4} \setminus \Lambda_3)$, that gives an m -tuple (w_1, w_2, \dots, w_m) fulfilling the Equations (1) and (2) with $w_{m+1} = w_1$.

4. Examples

Let us present the examples to support our conclusions.

4.1. Example

Consider the third order problem with $r = 1$,

$$\left. \begin{aligned}
 w_1'''(\varpi) + a_1(\varpi)f_1(w_2) &= 0, & \varpi \in [0, 1], \\
 w_2'''(\varpi) + a_2(\varpi)f_2(w_3) &= 0, & \varpi \in [0, 1], \\
 w_3'''(\varpi) + a_3(\varpi)f_3(w_1) &= 0, & \varpi \in [0, 1],
 \end{aligned} \right\} \tag{15}$$

$$\left. \begin{aligned}
 w_1(0) = 0, \quad w_1'(0) = 0, \quad w_1'(1) - 1 \int_0^1 w_1'(\xi)d\xi &= \beta_1, \\
 w_2(0) = 0, \quad w_2'(0) = 0, \quad w_2'(1) - \frac{1}{2} \int_0^1 \xi w_2'(\xi)d\xi &= \beta_2, \\
 w_3(0) = 0, \quad w_3'(0) = 0, \quad w_3'(1) - \frac{1}{3} \int_0^1 \xi^2 w_3'(\xi)d\xi &= \beta_3,
 \end{aligned} \right\} \tag{16}$$

where

$$a_1(\varpi) = \frac{1}{4}, a_2(\varpi) = \frac{1}{2}, a_3(\varpi) = \frac{3}{4}, g_1(\xi) = 1, \quad g_2(\xi) = \xi, \quad g_3(\xi) = \xi^2,$$

$$f_1(w_2) = w_2^2(1 + e^{-3w_2}), f_2(w_3) = w_3^2(1 - 3e^{-2w_3}), f_3(w_1) = w_1^2(1 + 4e^{-w_1}).$$

Then $f_{i0} = 0$ and $f_{i\infty} = \infty$ for $i = 1, 2, 3$. So, all the assumptions of Theorem 3.2 are met, and hence, the problem (15)-(16) has at least one positive solution by selecting β_1, β_2 and β_3 that are sufficiently small.

4.2. Example

Consider the third order problem with $r = 1$,

$$\left. \begin{aligned} w_1'''(\mathfrak{x}) + a_1(\mathfrak{x})f_1(w_2) &= 0, \mathfrak{x} \in [0, 1], \\ w_2'''(\mathfrak{x}) + a_2(\mathfrak{x})f_2(w_3) &= 0, \mathfrak{x} \in [0, 1], \\ w_3'''(\mathfrak{x}) + a_3(\mathfrak{x})f_3(w_1) &= 0, \mathfrak{x} \in [0, 1], \end{aligned} \right\} \quad (17)$$

$$\left. \begin{aligned} w_1(0) = 0, \quad w_1'(0) = 0, \quad w_1'(1) - 1 \int_0^1 w_1'(\xi)d\xi &= \beta_1, \\ w_2(0) = 0, \quad w_2'(0) = 0, \quad w_2'(1) - 2 \int_0^1 \xi w_2'(\xi)d\xi &= \beta_2, \\ w_3(0) = 0, \quad w_3'(0) = 0, \quad w_3'(1) - 3 \int_0^1 \xi^2 w_3'(\xi)d\xi &= \beta_3, \end{aligned} \right\} \quad (18)$$

where

$$\begin{aligned} a_1(\mathfrak{x}) = \mathfrak{x}, a_2(\mathfrak{x}) = \mathfrak{x}^2, a_3(\mathfrak{x}) = \mathfrak{x}^3, \quad g_1(\xi) = 1, \quad g_2(\xi) = \xi, \quad g_3(\xi) = \xi^2, \\ f_1(w_2) = w_2^{\frac{2}{3}}, f_2(w_3) = w_3^{\frac{3}{4}}, \quad f_3(w_1) = w_1^{\frac{1}{2}}. \end{aligned}$$

Then $f_{i0} = \infty$ and $f_{i\infty} = 0$ for $i = 1, 2, 3$. So, all the assumptions of Theorem 3.3 are met, and hence, the problem (17) and (18) has at least one positive solution by selecting β_1, β_2 and β_3 that are small enough.

5. Conclusion

This study employs the Guo-Krasnosel'skii fixed point theorem as a central tool to investigate the existence of positive solutions for an iterative system of higher order boundary value problems associated with non-homogeneous integral boundary conditions. This theorem, foundational in nonlinear analysis, provides conditions under which a compact operator has at least one fixed point within a cone in a Banach space. By applying this theorem, the study aims to rigorously establish criteria that guarantee the existence of positive solutions for the boundary value problems under consideration. It may be interesting that the researchers extend the results to multipoint boundary value problems and obtain multiple positive solutions by using various new fixed-point theorems.

Author contributions: Conceptualization, RPK and SN; methodology, RPK, SN and KN; validation, RPK, SN and KN; investigation, RPK, SN and KN; resources, RPK, SN and KN; writing—original draft preparation, RPK, SN and KN; writing—review and editing, RPK, SN and KN; supervision, SN. All authors have read and agreed to the published version of the manuscript.

Acknowledgments: The authors express their sincere gratitude to the referees for their insightful recommendations and suggestions.

Conflict of interest: The authors declare no conflict of interest.

References

1. Cannon JR. The solution of the heat equation subject to the specification of energy. *Quarterly of Applied Mathematics*. 1963; 21(2): 155–160. doi: 10.1090/qam/160437
2. Ionkin NI. Solution of a boundary value problem in heat conduction with a nonclassical boundary condition. *Differential Equations*. 1977; 13: 204–211.
3. Chegis RY. Numerical solution of the heat conduction problem with an integral condition. *Lithuanian Mathematical Proceedings*. 1984; 24: 209–215.
4. Dang QA, Dang QL. Existence results and iterative method for fully third order nonlinear integral boundary value problems. *Applications of Mathematics*. 2021; 66(5): 657–672. doi: 10.21136/am.2021.0040-20
5. Lv X, Wang L, Pei M. Monotone positive solution of a fourth-order BVP with integral boundary conditions. *Boundary Value Problems*. 2015; 2015(1). doi: 10.1186/s13661-015-0441-2
6. Benaicha S, Haddouchi F. Positive solutions of a nonlinear fourth-order integral boundary value problem. *Annals of West University of Timisoara-Mathematics and Computer Science*. 2016; 54(1): 73–86. doi: 10.1515/awutm-2016-0005 <https://doi.org/10.1515/awutm-2016-0005>
7. Liu K, Yang Y, Yang Y. Positive solutions to a BVP with two integral boundary conditions. *Annals of Applied Mathematics*. 2020; 36(3): 248–256.
8. Karaca IY, Fen FT. Positive Solutions of Nth-Order Boundary Value Problems with Integral Boundary Conditions. *Mathematical Modelling and Analysis*. 2015; 20(2): 188–204. doi: 10.3846/13926292.2015.1020531
9. Sreedhar N, Prasad KR, Balakrishna S. Existence of symmetric positive solutions for Lidstone type integral boundary value problems. *TWMS Journal of Applied and Engineering Mathematics*. 2018; 8(1a): 295–305.
10. Kanakayya N, Sreedhar N, Prasad KR. Existence and nonexistence results for higher order differential equations with non-homogenous integral boundary conditions. *Poincare Journal of Analysis & Applications*. 2021; 8(2): 119–131. doi:10.46753/pjaa.2021.v08i02.001
11. Henderson J, Ntouyas SK, Purnaras IK. Positive solutions for systems of second order four-point nonlinear boundary value problems. *Communications in Applied Analysis*. 2008; 12(1): 29–40.
12. Henderson J, Ntouyas SK, Purnaras IK. Positive solutions for systems of generalized three-point nonlinear boundary value problems. *Commentationes Mathematicae Universitatis Carolinae*. 2008; 49(1): 79–91.
13. Prasad KR, Sreedhar N, Kumar KR. Solvability of iterative systems of three-point boundary value problems. *TWMS Journal of Applied and Engineering Mathematics*. 2013; 3(2): 147–159.
14. Oz D, Karaca IY. Eigenvalue intervals for iterative systems of second-order nonlinear equations with impulses and m-point boundary conditions. *Fixed Point Theory*. 2024; 25(1): 289–308. doi: 10.24193/fpt-ro.2024.1.18
15. Bouteraa N, Benaicha S, Djourdem H. Positive Solutions for Systems of Fourth Order Two-Point Boundary Value Problems with Parameter. *Journal of Mathematical Sciences and Modelling*. 2019; 2(1): 30–38. doi: 10.33187/jmsm.432678
16. Henderson J, Ntouyas S. Positive solutions for systems of nth order three-point nonlocal boundary value problems. *Electronic Journal of Qualitative Theory of Differential Equations*. 2007; (18): 1–12. doi: 10.14232/ejqtde.2007.1.18
17. Prasad KR, Rashmita M, Sreedhar N. Solvability of higher order three-point iterative systems. *Ufimsky Journal of Mathematics*. 2020; 12(3): 107–122. doi: 10.13108/2020-12-3-107
18. Namburi S, Namana K, Rajendra PK. Solvability of Higher Order Iterative System with Non-Homogeneous Integral Boundary Conditions. *Contemporary Mathematics*. 2022. doi: 10.37256/cm.3220221300
19. Mansouri B, Ardjouni A, Djoudi A. Positive solutions of nonlinear fourth order iterative differential equations with two-point and integral boundary conditions. *Nonautonomous Dynamical Systems*. 2021; 8(1): 297–306. doi: 10.1515/msds-2020-0139
20. Chouaf S, Khemis R, Bouakkaz A. Some existence results on positive solutions for an iterative second-order boundary-value problem with integral boundary conditions. *Bulletin of the Paraná Mathematical Society*. 2022; 40: 1–10. doi: 10.5269/bspm.52461
21. Henderson J, Luca R, Tudorache A. Positive Solutions for a System of Second-Order Differential Equations with Integral Boundary Conditions. *International Journal of Applied Physics and Mathematics*. 2016; 6(3): 138–149. doi: 10.17706/ijapm.2016.6.3.138-149

22. Sun JP, Yang XM, Zhao YH. Existence of Positive Solution to System of Nonlinear Third-Order Three-Point BVPs. *Journal of Function Spaces*. 2016; 2016: 1–6. doi: 10.1155/2016/6874643
23. Eloe PW, Henderson J. Positive solutions for higher order ordinary differential equations. *Electronic Journal of Differential Equations*. 1995; 1995(3): 1–8. https://doi.org/10.1142/9789812831118_0001
24. Sun JP, Li HB. Monotone Positive Solution of Nonlinear Third-Order BVP with Integral Boundary Conditions. *Boundary Value Problems*. 2010; 2010(1): 874959. doi: 10.1155/2010/874959
25. Mahdy AMS. Stability, existence, and uniqueness for solving fractional glioblastoma multiforme using a Caputo–Fabrizio derivative. *Mathematical Methods in the Applied Sciences*. 2023. doi: 10.1002/mma.9038
26. Mahdy AMS. Numerical Solution and Optimal Control for Fractional Tumor Immune Model. *Journal of Applied Analysis & Computation*. 2024; 14(5): 3033–3045. doi: 10.11948/20240053
27. El-Mesady A, Elsadany AA, Mahdy AMS, et al. Nonlinear dynamics and optimal control strategies of a novel fractional-order lumpy skin disease model. *Journal of Computational Science*. 2024; 79: 102286. doi: 10.1016/j.jocs.2024.102286
28. Mohammed D Sh, Abdou MA, Mahdy AMS. Dynamical investigation and numerical modeling of a fractional mixed nonlinear partial integro-differential problem in time and space. *Journal of Applied Analysis & Computation*. 2024; 14(6): 3458–3479. <http://dx.doi.org/10.11948/20240050>
29. Mahdy AMS, Abdou MA, Mohamed DS. Numerical solution, convergence and stability of error to solve quadratic mixed integral equation. *Journal of Applied Mathematics and Computing*. 2024. doi: 10.1007/s12190-024-02194-1
30. Bueckner HF. Positive Solutions of Operator Equations (M. A. Krasnoselskii). *SIAM Review*. 1966; 8(1). doi: 10.1137/1008029
31. Guo D, Lakshmikantham V. *Nonlinear Problems in Abstract Cones*. Academic Press; 1988.
32. Munkres JR. *Topology: A First Course*. Pearson College Div; 1974.