

Article

A study on membrane enzyme $\text{Na}^+\text{-K}^+\text{-ATPase}$ in lindane exposed fish, *Channa punctatus*

Aradhna Gupta, Bechan Sharma*

Department of Biochemistry, University of Allahabad, Allahabad 211002, India

* Corresponding author: Bechan Sharma, sharmabi@yahoo.com

CITATION

Gupta A, Sharma B. A study on membrane enzyme $\text{Na}^+\text{-K}^+\text{-ATPase}$ in lindane exposed fish, *Channa punctatus*. Journal of Toxicological Studies. 2024; 2(2): 1238.
<https://doi.org/10.59400/jts.v2i2.1238>

ARTICLE INFO

Received: 26 April 2024

Accepted: 3 June 2024

Available online: 23 July 2024

COPYRIGHT



Copyright © 2024 author(s).

Journal of Toxicological Studies is published by Academic Publishing Pte. Ltd. This work is licensed under the Creative Commons Attribution (CC BY) license.

<https://creativecommons.org/licenses/by/4.0/>

Abstract: $\text{Na}^+\text{-K}^+\text{-ATPase}$ is a membrane bound enzyme responsible for the transport of ions through the membrane and immediate release of energy. This enzyme is known to be an early target for oxygen radical induced damage to intact cell. Exposure of *C. punctatus* to subacute concentrations of lindane for 96 h caused significant reduction in the activities of $\text{Na}^+\text{-K}^+\text{-ATPase}$ in all the tissues of the fish tested; brain being maximally affected and the heart being least affected organ at the highest concentration of lindane (0.1 mg/L). The effect of pesticide was concentration dependent. The percent decrease in the activity of $\text{Na}^+\text{-K}^+\text{-ATPase}$ in brain, gills, heart, kidney, liver and muscle was found to be 36.7, 23.4, 19.2, 29, 22.9 and 29.7, respectively. The order of level of enzyme activity recorded was as following: liver > gills > kidney > brain > muscle > heart in the control.

Keywords: Lindane; Na-K-ATPase; organs

1. Introduction

The aquatic ecosystem is an open system exposed to all different kind of pollutants, toxicants, surfactants thus polluting the aquatic environment. Their direct discharge without any pre-treatment either leads to large scale destruction of aquatic life or accumulation in water, soil/bioaccumulation in biotic aquata. Though, the pollutants could be biodegradable or non-biodegradable, these pollutants sometimes, decrease the rate of decay of biodegradables. Thus, the increase in the contaminants may allow their environmental persistence for longer period. Excessive use of nitrate and phosphate fertilizers may lead to eutrophication, thereby reducing the amount of oxygen in the aquatic biota and increasing the biological oxygen demand of that water for prolonged period. Industrial smokes, burning of woods, petroleum, and vehicles fumes all gave rise to gases like sulphur dioxide, nitrogen oxides, and carbon dioxide; lead, and the particulate matters, all have caused serious harm to the environment and humans [1–3]. Reports on the increased levels of heavy metals and pesticides and their effects on oxidative stress have been exhaustively documented [4–8]. Many bacteria like *Vibrio anguillarum*, *Aeromonas*, *Flavobacterium*, *Pseudomonas*, *Serratia*, and *Yersina*, etc. have been shown to grow in water, which has less oxygen, increased organic matter, and unsuitable pH for the aquatic life.

The presence of organic matter may be due to leakage from septic tanks or contamination by domestic sewage [9]. Aquatic life may suffer from diseases like fin rot, papilloma, hyper-neoplasia, and gill diseases etc. Gills are the main respiratory organs of fish; it regulates ion concentration and osmotic balance for survival in unfavorable concentrations [10]. Decrease in function of gills has been reported when exposed to pH of water less or more than 7 [11]. Lindane is an organochlorine,

hydrophobic, and highly persistent pesticide. Due to its lipophilic nature, it gets easily bioaccumulated in aquatic organisms. Lindane is strongly adsorbed on soils that contain a large amount of organic matter. It can move downward by capillary action through the soil with water from rainfall or artificial irrigation. In the UV light it undergoes rapid dichlorination or degradation to form pentachloro cyclohexenes and tetrachlorocyclohexenes. The fish are able to bioaccumulate due to direct exposure to chemicals in water, and ingestion of contaminated food or preys [12]. Their accumulation in low concentration in aquatic animals generate warning signals about the environment. Ultimately in long run when these fish are eaten by humans or animals, the pesticide residues accumulate in different organs of the exposed living systems and hence may pose serious health problems [13]. The pesticide and its residues are known to get strongly adsorbed on clays and sediments of surface water where several fish population reside for feeding.

Sodium Potassium ATPases ($\text{Na}^+\text{-K}^+\text{-ATPase}$, EC 3.6. 3.9) is a membrane bound sulfhydryl containing oligomeric enzyme whose function is critical for the maintenance of cell viability. It works as an electrogenic P-type pump involved in monovalent ion transport across membranes utilizing energy of ATP hydrolysis. It pumps 3 sodium outside (extracellular) and 2 potassium inside the cell membrane (intracellular) for 1 ATP. The concentration gradient created due to ions transport across the membranes help in other secondary transports [14,15]. This pump has many diverse functions like contractions, signalling, homeostasis (osmoregulation), and cell-cell adhesions [16,17]. It also governs many physiological processes like reabsorption, filtration, pH, electrolytes osmotic regulation by kidneys [18], sperm motility [19], and action potential in neurons [20–23]. This pump consists of 3 subunits α , β and FXYD. The α subunit is for ion transport (catalytic) and is under influence of β (auxiliary) and FXYD (tissue specific) subunit [24]. The $\beta 2$ subunit of $\text{Na}^+\text{-K}^+\text{-ATPase}$ is responsible for regulation of egg development in *Aedes aegypti*, the causative organism of many vectors borne diseases like yellow fever, zika, dengue, chicken guinea. The reduced expression of mRNA of $\beta 2$ $\text{Na}^+\text{-K}^+\text{-ATPase}$ in the knockdown mice showed less egg formation, which directly reduced their population and so less spread of disease [25]. In humans, decreased heart functions or heart failures have been reported due to decrease in its activity [26]. In plants, it helps in nutrient uptake, root development, stomatal regulation, and response to environmental stresses. It helps plants to maintain proper ion balance, participate in nutrient absorption, and regulate turgor pressure in cell. In the environment, it acts as marker for organisms facing challenges such as varying salinity, ion concentrations, or osmotic stress. In the present study, the effect of subacute concentrations of lindane (0.025, 0.05, 0.1 mg/L) on the activity of $\text{Na}^+\text{-K}^+\text{-ATPase}$ was assessed in order to evaluate the perturbations in the transport of ions in different fish tissues exposed for 96 h.

2. Materials and methods

2.1. Experimental design and exposure to lindane

The healthy and active fish of 30–40 gm with no signs of any diseases or external injury were equally distributed in four aquaria of 1×1 ft. The subacute concentrations

of lindane (0.025, 0.05, 0.1 mg/L) prepared in acetone were used for the exposure of *C. punctatus* for 96 h. In control group, equal volume of acetone was added. All aquaria were constantly aerated during the period of exposure by aerator and the fish were fed properly. The water of all 4 aquaria was changed in 24 h and replenished with fresh lindane.

2.2. Preparation of cell free extracts and biochemical assays

- 1) Protein determination: The quantitative estimation of total protein in various tissue extracts and solutions were done according to the known procedure [27]. The samples were homogenized 10% (w/v) in 0.05M sodium phosphate buffer, pH 7.4 and centrifuged at 10,000 rpm for 10min in cold (4 °C) condition. The supernatants were collected in labelled vials and volume was noted. Determination of protein was done using Folin-Ciocalteu reagent. The bovine serum albumin (BSA) was used as a standard. A blank was prepared, which contained all reagents but no protein. The intensity of blue color was measured colorimetrically at 620 nm.
- 2) Assay of Na⁺-K⁺-ATPase activity: The Na⁺-K⁺-ATPase activity was determined as inorganic phosphorus (Pi) production using the method of Svobaca and Mossinger [28] and Fiske and Subbarow [29]. The reaction mixture is given in **Table 1**.

All reagents were obtained from Sigma Chemical Company, U.S.A. The pesticide used in the experiments was from Rallis India limited. The double distilled water was used for all biochemical experiments.

Table 1. Process for assay of Na⁺-K⁺-ATPase activity.

10% (w/v) tissue homogenized in 0.25 M sucrose (pH 7.4)	
15 min centrifugation 12,000 rpm at 4 °C	
Supernatant collected	
1	2
0.2 mL of 200 mM KCl	
0.2 mL of 1 M NaCl	0.1 mL of 1000 mM MgCl ₂
0.1 mL of 1000 mM MgCl ₂	1.0 mL of 200 mM Tris buffer, pH 7.4
1.0 mL of 200 mM Tris buffer, pH 7.4	100–200 µg supernatant
0.2 mL distilled water	0.16 mL distilled water
100–200 µg supernatant	0.2 mL of 10 mM ouabain
Leave 5 min/35–37 °C	
0.2 mL of 25 mM ATP (di sodium salt)	0.2 mL of NaCl + 0.2 mL of 25 mM ATP
Leave 15 min/35–37 °C	
1 mL of 10% TCA	
3000 rpm/5 min	
Supernatant collected	
0.5 mL supernatant	
3.0 mL DW	
0.5 mL of 2.5% ammonium molybdate in 5N H ₂ SO ₄	

Table 1. (Continued).

10% (w/v) tissue homogenized in 0.25 m sucrose (pH 7.4)	
15 min centrifugation 12,000 rpm at 4 °C	
Supernatant collected	
1	2
0.2 mL of 1, 2, 3, 4 ANSA	
Vortexed/10 min	
Reading at 600 nm	
Difference between 1 and 2 gives activity of Na ⁺ -K ⁺ -ATPase in μmol Pi liberated/mg protein/h.	
ANSA = aminonaphthol sulphonic.	

2.3. Statistical analysis

The values were presented as means ± standard error of mean (SEM) of observed data of three to five replicates. Graph pad prism version 3.0 (GraphPad Prism Software Inc., San Diego, CA, USA) was used to analysis the data. Results obtained from treated and control fish were compared using Turkeys H test.

3. Results and discussion

Effect of lindane on the activity of Na⁺-K⁺-ATPase in different organs of lindane exposed fish

The data demonstrated highest activity of Na⁺-K⁺-ATPase to be present in liver (41.67 ± 0.35 units/mg protein) and lowest in heart (16.73 ± 0.11 units/mg protein) of the control fish. The activities of Na⁺-K⁺-ATPase in other fish tissues such as gills, kidney, brain and muscle were recorded as 33.78 ± 0.23, 32.87 ± 0.19, 30.29 ± 0.07 and 21.29 ± 0.09 units/mg protein, respectively. The order of level of enzyme activity recorded in the control fish tissues was as following: liver > gills > kidney > brain > muscle > heart (Table 2, Figure 1).

Table 2. Effect of lindane (mg/L) on the specific activity of Na⁺-K⁺-ATPase (units/mg protein) in different tissues of *C. punctatus* exposed for 96 h.

Lindane	Gills	Heart	Kidney	Liver	Muscle	Brain
0	33.78 ± 0.23	16.73 ± 0.11	32.87 ± 0.19	41.67 ± 0.35	21.29 ± 0.09	30.29 ± 0.07
0.025	30.59 ± 0.21 (-9.42)	15.69 ± 0.09 (-6.21)	29.79 ± 0.18 (-9.37)	39.27 ± 0.33 (-5.76)	19.17 ± 0.08 (-9.95)	25.31 ± 0.05 (-16.44)
0.05	28.67 ± 0.22 (-14.98)	14.87 ± 0.07 (-11.10)	25.66 ± 0.15 (-21.97)	37.09 ± 0.32 (-11.01)	17.15 ± 0.07 (-19.47)	22.11 ± 0.06 (-27.02)
0.1	25.87 ± 0.20 (-23.41)	13.52 ± 0.08 (-19.19)	23.35 ± 0.16 (-28.94)	32.13 ± 0.29 (-22.92)	14.97 ± 0.07 (-29.70)	19.17 ± 0.04 (-36.68)

Values are represented as μm of Pi released/h/mg wet weight of tissue. Each value represents the mean ± SEM of ten different observations. Values in parenthesis are percent change over control. The (-) sign represents decrease over control. *h* represents time in hour. SEM = standard error of mean.




Figure 1. Lindane exposure at different concentrations (0, 0.025, 0.05, 0.1 ppm) on the specific activity of Na⁺-K⁺-ATPase (units/mg protein) in different tissues of *C. punctatus* exposed for 96 h.

From the Turkey's HSD test all pairwise comparisons between different concentrations (0.025, 0.05, 0.1 mg/L) and control (0 mg/L) were statistically significant with $p < 0.05$. This indicates that there are significant differences in the mean values between the control and each of the concentrations, as well as between the different concentrations themselves for organs brain, muscle and liver. The non-significant differences were observed in the gills and kidney with 0.1 vs. 0.05 mg/L and heart 0.05 vs. 0.025 mg/L (**Table 3**).

Table 3. Statistical inference of data based on pairwise comparisons between different concentrations of lindane (mg/L).

Lindane	Gills	Heart	Kidney	Liver	Muscle	Brain
0.025 vs. 0	$P < 0.001$	S	$P < 0.001$	S	$P < 0.001$	S
0.05 vs. 0	$P < 0.001$	S	$P < 0.001$	S	$P < 0.001$	S
0.1 vs. 0	$P < 0.001$	S	$P < 0.001$	S	$P < 0.001$	S
0.05 vs. 0.025	$P < 0.001$	S	$P = 0.34$	NS	$P < 0.001$	S
0.1 vs. 0.025	$P < 0.001$	S	$P < 0.001$	S	$P < 0.001$	S
0.1 vs. 0.05	$P = 0.52$	NS	$P < 0.001$	S	$P = 0.16$	NS

S = Significant, NS = Not significant.

The treatment of the fish with three subacute concentrations of lindane displayed marked decrease in the activity of Na⁺-K⁺-ATPase in all the tissues tested i.e. brain being maximally affected at all of these concentrations. At lowest concentration of lindane (0.025 mg/L), the fish tissues such as gills, kidney and muscle showed same level of percent inhibition (about 9%–10%) in the enzyme activity (**Table 2**).

At the highest concentration of the pesticide (0.1 mg/L), the brain of the fish exhibited maximum decrease by 37% in the enzyme activity and the heart indicated minimum decrease by 19% after 96 h of treatment duration. The decrease in the

enzyme activity was in concentration dependent manner. The percent decrease in the activity of Na⁺-K⁺-ATPase in brain, gills, heart, kidney, liver and muscle was found to be 36.7, 23.4, 19.2, 29, 22.9 and 29.7, respectively. The data indicated that the gills and liver were affected by lindane to similar level showing about 23% decrease in enzyme activity. Further, the kidney and muscle displayed almost same level of enzyme inhibition to about 29% when the fish was treated with lindane at 0.1 mg/L concentration for 96h. The extent of inhibition in the enzyme activity from these fish tissues treated with the highest pesticide concentration (0.1 mg/L) was found in the following order: brain > muscle = kidney > gills = liver > heart (**Table 2**).

This enzyme is known to be an early target for oxygen radical induced damage to intact cell [30,31]. Sharma observed significant reduction of liver ATPase activity in *C. gaucha* upon endosulphan exposure [32]. Oruc et al. reported reduction of liver activity in *Tillapia zilli* and *O. Niloticus* and suggested that increased lipid peroxidation disturbed the anatomical integrity of the biomembrane and diminished its fluidity leading to inhibition of activities of several membrane bound enzymes including Na⁺-K⁺-ATPase [33]. Very recently, cypermethrin at lethal (5.03 µg/L) and sublethal (1.02 µg/L) concentrations has been shown to cause significant alterations in gills, liver and muscle of the fish *Cirrhinus mrigala* [34].

Similar observations have been reported in *C. punctatus* exposed to pyrethroids [35], *Clarias gariepinus* juvenile exposed to oxadiazon [35]. Increased activity in gills of silver cat fish (*Rhamdia quelen*) was reported in water pH 9.0 and no significant change in kidney [36]. In hypotonic environment, kidneys of fish species excrete more dilute urine to maintain homeostasis [37] and in hypertonic they suffer from dehydration and so they drink more sea water excreting out extra salts [38,39]. At lethal concentration, the insecticide caused increased in enzyme activity whereas at sublethal concentration the enzyme activity decreased [40]. Maiti exposed the fish *Clarius batrachus* to 5.69 mg/L and 11.38 mg/L of chromium (III) for 96h and reported decrease in activity in brain [41–45]. This inactivity can lead to diverse alterations in the neurons such as partial membrane depolarization, Ca⁺² influx, altered neurotransmitter release and even apoptosis and all can be co-related to functional deficits in the brain [46].

4. Conclusions

The significance of sodium-potassium ATPase (Na⁺-K⁺-ATPase) in the environment is its role played in maintaining cellular homeostasis both in plants and animals. It contributes to the adaptability and survival of organisms in diverse environmental conditions and serves as a sensitive indicator of environmental stress and toxicity. The exposure of the fish to varying subacute concentrations of lindane have caused significant perturbations in the activity of this enzyme in different organs of the fish tested. The results may serve as indicator of pesticide contamination in water and better management of pesticide application in agricultural practices to secure environmental health.

Author contributions: Conceptualization, methodology and drafting, BS; performed experiment, AG. All authors have read and agreed to the published version of the

manuscript.

Acknowledgments: The authors are grateful to University of Allahabad for providing facilities for carrying out the present work.

Conflict of interest: The authors declare no conflict of interest.

References

1. Gupta A, Rai DK, Pandey RS, et al. Analysis of some heavy metals in the riverine water, sediments and fish from river Ganges at Allahabad. *Environmental Monitoring and Assessment*. 2008; 157(1–4): 449–458. doi: 10.1007/s10661-008-0547-4
2. German AV, Zakonov VV, Mamontov AA. Organochlorine compounds in bottom sediments, benthos, and fish in the volga pool of the Rybinsk Reservoir. *Water Resources*. 2010; 37(1): 84–88. doi: 10.1134/s0097807810010082
3. Gupta MA. A Case Study on the Bioaccumulated Organochlorines in Fish Related to their inhabitant Res. *Rev. J Ecol. Environ*. 2021; 9(5).
4. Gupta A, Sharma B. Evaluation of Levels of Phosphatases in the Lindane Exposed Fish, *Channa punctatus*. *Journal of Biomedical Research & Environmental Sciences*. 2023; 4(3): 555–561. doi: 10.37871/jbres1710
5. Jackson DA, Gardner DR. The effects of some organochlorine pesticide analogs on salmonid brain ATPases. *Pesticide Biochemistry and Physiology*. 1973; 2(4): 377–382. doi:10.1016/0048-3575(73)90049-7
6. Gupta A, Siddiqi NJ, Sharma B. Bioaccumulation and Biochemical Studies of Toxicants in Fish on AChE. *Open J Pathol Toxicol Res*. 2021; 1(1).
7. Gupta A, Sharma B. Acute Chronic Toxicity of Lindane in *Channa punctatus*. *Open J Pathol Toxicol Res*. 2021; 1(1).
8. Gupta A. An Evaluation of Lactate Dehydrogenase in the Lindane. *International Journal of Animal Biotechnology and Applications*. 2021; 7(1): 2455–7315.
9. Austin B. The effects of pollution on fish health. *Journal of Applied Microbiology*. 1998; 85(S1): 234S–242S. doi: 10.1111/j.1365-2672.1998.tb05303.x
10. Zimmer AM, Perry SF. Physiology and aquaculture: A review of ion and acid-base regulation by the gills of fishes. *Fish and Fisheries*. 2022; 23(4): 874–898. doi: 10.1111/faf.12659
11. Garcia Parra J, Baldisserotto B. Effect of Water pH and Hardness on Survival and Growth of Freshwater Teleosts. In: *Fish Osmoregulation*. Boca Raton; 2007. pp. 135–150. doi: 10.1201/b10994-6
12. Almeida CL, Aguiar LH, Moraes G. Effect of methyl parathion on the muscle and brain acetylcholinesterase activity of matrinxã (*Brycon cephalus*). *Ciência Rural*. 2005; 35(6): 1412–1416. doi: 10.1590/s0103-84782005000600029
13. Metcalf RL. *Pesticides in Aquatic Environments*. Springer US; 1977. doi: 10.1007/978-1-4684-2868-1
14. Lucu Ć, Towle DW. Na⁺/K⁺-ATPase in gills of aquatic crustacea. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*. 2003; 135(2): 195–214. doi: 10.1016/s1095-6433(03)00064-3
15. Thorsen K, Drengstig T, Ruoff P. Transepithelial glucose transport and Na⁺/K⁺ homeostasis in enterocytes: An integrative model. *American Journal of Physiology-Cell Physiology*. 2014; 307(4): C320–C337. doi: 10.1152/ajpcell.00068.2013
16. Nie Y, Bai F, Chaudhry MA, et al. The Na/K-ATPase α 1 and c-Src form signaling complex under native condition: A crosslinking approach. *Scientific Reports*. 2020; 10(1). doi: 10.1038/s41598-020-61920-4
17. Pivovarov AS, Calahorra F, Walker RJ. Na⁺/K⁺-pump and neurotransmitter membrane receptors. *Invertebrate Neuroscience*. 2018; 19(1). doi: 10.1007/s10158-018-0221-7
18. Mernissi G, Barlet-Bas C, Khadouri C, et al. Characterization and localization of ouabain-insensitive Na-dependent ATPase activities along the rat nephron. *Biochim Biophys Acta*. 1991; 1064(2): 205–211. doi: 10.1016/0005-2736(91)90303-P
19. Jimenez T, McDermott JP, Sánchez G, et al. Na/K-ATPase α 4 isoform is essential for sperm fertility. *Proceedings of the National Academy of Sciences*. 2010; 108(2): 644–649. doi: 10.1073/pnas.1016902108
20. Clausen MV, Hilbers F, Poulsen H. The Structure and Function of the Na/K-ATPase Isoforms in Health and Disease. *Frontiers in Physiology*. 2017; 8. doi: 10.3389/fphys.2017.00371
21. Attwell D, Laughlin SB. An Energy Budget for Signaling in the Grey Matter of the Brain. *Journal of Cerebral Blood Flow & Metabolism*. 2001; 21(10): 1133–1145. doi: 10.1097/00004647-200110000-00001
22. Lei J, Nowbar S, Mariash CN, et al. Thyroid hormone stimulates Na-K-ATPase activity and its plasma membrane insertion

- in rat alveolar epithelial cells. *American Journal of Physiology-Lung Cellular and Molecular Physiology*. 2003; 285(3): L762–L772. doi: 10.1152/ajplung.00376.2002
23. Pirahanchi Y, Jessu R, Aeddula NR. Physiology, Sodium Potassium Pump. Available online: <https://www.ncbi.nlm.nih.gov/books/NBK537088/> (accessed on 13 March 2023).
 24. Mercer RW, Biemesderfer D, Bliss DP, et al. Molecular cloning and immunological characterization of the gamma polypeptide, a small protein associated with the Na/K-ATPase. *The Journal of cell biology*. 1993; 121(3): 579–586. doi: 10.1083/jcb.121.3.579
 25. Martinez NP, Pinch M, Kandel Y, et al. Knockdown of the Sodium/Potassium ATPase Subunit Beta 2 Reduces Egg Production in the Dengue Vector, *Aedes aegypti*. *Insects*. 2023; 14(1): 50. doi: 10.3390/insects14010050
 26. Kjeldsen K. Myocardial Na/K-ATPase: Clinical aspects. *Exp Clin Cardiol*. 2003; 8(3): 131–133.
 27. Lowry OH, Rosebrough NJ, Farr AI, Randall RJ. Protein measurement with Folin-Phenol reagent. *J. Biol. Chem.* 1951; 193: 265–275. doi: 10.1016/S0021-9258(19)52451-6
 28. Svobaca P, Mossinger B. Catecholamine and brain microsomal Na⁺/K⁺-ATPase-1, protection against lipoperoxidative damages. *Biochem. Pharmacol.* 1981; 30: 427–432. doi: 10.1016/0006-2952(81)90626-2
 29. Fiske CH, Subbarow Y. Colourimetric determination of phosphorous. *J Biol. Chem.* 1925; 66: 375–400. doi: 10.1016/S0021-9258(18)84756-1
 30. Kim MS, Akera T. O₂ free radicals: cause of ischemia-reperfusion injury to cardiac Na⁺-K⁺-ATPase. *American Journal of Physiology-Heart and Circulatory Physiology*. 1987; 252(2): H252–H257. doi: 10.1152/ajpheart.1987.252.2.h252
 31. Kako K, Kato M, Matsuoka T, et al. Depression of membrane-bound Na⁺-K⁺-ATPase activity induced by free radicals and by ischemia of kidney. *American Journal of Physiology-Cell Physiology*. 1988; 254(2): C330–C337. doi: 10.1152/ajpcell.1988.254.2.c330
 32. Sharma RM. Effect of endosulfan on adenosine triphosphatase (ATPase) activity in liver, kidney, and muscles of *Channa gachua*. *Bulletin of Environmental Contamination and Toxicology*. 1988; 41(3): 317–323. doi: 10.1007/bf01688873
 33. Ozcan Oruc E, Uner N, Tamer L. Comparison of Na⁺-K⁺-ATPase Activities and Malondialdehyde Contents in Liver Tissue for Three Fish Species Exposed to Azinphosmethyl. *Bulletin of Environmental Contamination and Toxicology*. 2002; 69(2): 271–277. doi: 10.1007/s00128-002-0057-y
 34. Prashanth MS, David M. Impact of Cypermethrin on Na⁺-K⁺, Ca²⁺ and Mg²⁺ ATPases in Indian Major Carp, *Cirrhinus mrigala* (Hamilton). *Bulletin of Environmental Contamination and Toxicology*. 2009; 84(1): 80–84. doi: 10.1007/s00128-009-9864-8
 35. Kumar A, Sharma B, and Pandey RS. Toxicological assessment of the pyrethroids insecticides with special reference to cypermethrin and λ-cyhalothrin in fresh water fishes. *Int. J. Biol. Med Res.* 2010; 1(4): 315–325.
 36. Oluah NS, Mgbenka BO, Nwani CD, et al. Tissue-specific changes in Ca²⁺-ATPase and Na⁺/K⁺-ATPase activities in freshwater African catfish *Clarias gariepinus* juvenile exposed to oxadiazon. *The Journal of Basic and Applied Zoology*. 2020; 81(1). doi: 10.1186/s41936-020-00186-8
 37. Marx MTS, Souza C de F, Almeida APG, et al. Expression of Ion Transporters and Na⁺/K⁺-ATPase and H⁺-ATPase Activities in the Gills and Kidney of Silver Catfish (*Rhamdia quelen*) Exposed to Different pHs. *Fishes*. 2022; 7(5): 261. doi: 10.3390/fishes7050261
 38. Esbaugh AJ, Brix KV, Grosell M. Na⁺/K⁺-ATPase isoform switching in zebrafish during transition to dilute freshwater habitats. *Proceedings of the Royal Society B: Biological Sciences*. 2019; 286(1903): 20190630. doi: 10.1098/rspb.2019.0630
 39. McCormick SD, Regish AM, Christensen AK. Distinct freshwater and seawater isoforms of Na⁺/K⁺-ATPase in gill chloride cells of Atlantic salmon. *Journal of Experimental Biology*. 2009; 212(24): 3994–4001. doi: 10.1242/jeb.037275
 40. Yang WK, Hsu AD, Kang CK, et al. Intestinal FXD12 and sodium-potassium ATPase: A comparative study on two euryhaline medakas in response to salinity changes. *PLOS ONE*. 2018; 13(7): e0201252. doi: 10.1371/journal.pone.0201252
 41. Maiti AK, Paul G, Maity B, et al. Chromium III Exposure Inhibits Brain Na⁺/K⁺-ATPase Activity of *Clarias batrachus* L. Involving Lipid Peroxidation and Deficient Mitochondrial Electron Transport Chain Activity. *Bulletin of Environmental Contamination and Toxicology*. 2009; 83(4): 479–483. doi: 10.1007/s00128-009-9827-0
 42. Gupta A, Sharma B. A study on Transaminases in Lindane Exposed Fish *C. punctatus*. *Journal of Biomedical Research & Environmental Sciences*. 2023; 4(6): 1100–1107. doi: 10.37871/jbres1773
 43. Gupta A, Sharma B. Oxidative stress biomarkers in a living cell. *MOJ Toxicol*. 2023; 7(1): 38–43. doi: 10.15406/mojt.2023.07.00176

44. Gupta A. A Study on Analysis of Water Quality of Two Rivers Ganges and Yamuna. *J Waste Manage Xenobio*. 2023; 6(3): 000191.
45. Gupta A, Sharma B. A review on water pollution by γ HCH (lindane) and its removal using nanomaterials. *Journal of Toxicological Studies*. 2023; 1(1): 195. doi: 10.59400/jts.v1i1.195
46. Lijnen P, Hespel P, Lommelen G, et al. Intracellular sodium, potassium and magnesium concentration, ouabain sensitive rubidium uptake and sodium efflux and Na^+/K^+ cotransport activity in erythrocytes of normal male subjects studied on two occasions. *Methods Find Exp Clin Pharmacol*. 1986; 8: 525–533.