

Toxicological evaluation of Ultrapure and Potent Tannic Acid (UPPTA) by inhalation exposure

Lu-Ping Lu^{1,2}, Yi-Wen Mao¹, Theodore Tsai¹, Yi-An Lai¹, Teh-Ying Chou^{2,3,5}, Guochuan Emil Tsai^{1,2,4,*}

¹ Department of Research and Development, SyneuRx International (Taiwan) Corp., New Taipei 221416, Taiwan

² Institute of Biochemistry and Molecular Biology, National Yang Ming Chiao Tung University, Taipei 112304, Taiwan

³ Graduate Institute of Clinical Medicine, Taipei Medical University, Taipei 11031, Taiwan

⁴ Department of Psychiatry and Biobehavioral Science, UCLA School of Medicine, Los Angeles 90024, California, USA

⁵ Department of Pathology and Precision Medicine Research Center, Taipei Medical University Hospital, Taipei Medical University, Taipei 112304, Taiwan

* Corresponding author: Guochuan Emil Tsai, tsaimdphd@ucla.edu

ARTICLE INFO

Received: 16 October 2023

Accepted: 19 December 2023

Available online: 26 December 2023

doi: 10.59400/jts.v1i1.245

Copyright © 2023 Author(s).

Journal of Toxicological Studies is published by Academic Publishing Pte. Ltd. This article is licensed under the Creative Commons Attribution License (CC BY 4.0).

<https://creativecommons.org/licenses/by/4.0/>

ABSTRACT: The therapeutic options for coronal virus infections are limited. As SARS-CoV-2 directly targets the lungs and causes lung damage, treatment of COVID-19 with inhalants may offer more advantages over oral administration. Inhaled drug delivery provides a higher drug concentration in the target organ, where SARS-CoV-2 proliferates. In this study, we evaluated the potential systemic toxicity, relevant target organ toxicity, and toxicokinetics of Airneclu[®], Ultrapure, and Potent Tannic Acid (UPPTA) by metered-dose inhaler (MDI) inhalation to rodent and canine species once a day for 2 consecutive weeks. We further investigated the reversibility of the toxicity following a 3-week recovery period. No mortality related to the test article was observed in all the dose groups. Neither abnormalities related to the test article nor toxicologically significant changes were observed in both rodent and canine studies. In pathological examination, alveolar macrophage aggregation, perivascular/interstitial/alveolar inflammatory cell infiltration, and alveolar/bronchial epithelium hyperplasia were noted in the lung with bronchial involvement. However, after a 3-week recovery period, a substantial recovery was observed. There is limited systemic exposure to the inhalation administration. Therefore, inhalation of Airneclu[®] UPPTA is safe to administer for respiratory disorders like COVID-19.

KEYWORDS: COVID-19; SARS-CoV-2; metered-dose inhaler; inhaled toxicity; pulmonary pathology

1. Introduction

Tannic acids, a group of tannins enriched in several species of sumac and Aleppo oak, are naturally occurring hydrolysable polyphenol compounds with galloyl moieties expanding from a central glucose moiety. The structures of tannic acids are composed of 1 to 10 units of galloyl moieties per molecule. Throughout human history, tannic acids have been used as (part of) herbal medicine worldwide for thousands of years. For example, tannic acids have been used to treat diarrhea, bleeding, coughing, and detoxification in China^[1].

The study of the anti-viral activity of tannic acids can be traced back more than 70 years. Green first reported that tannic acids can inhibit influenza A virus multiplication and hemagglutination^[2]. More recently, tannic acids extracted from tea showed inhibitory activity against SARS-CoV with an IC₅₀ of 3 μM^[3]. The main protease of coronaviruses is essential for the processing of the replicase complex nonstructural proteins. In addition, the nasal epithelium is critical in the processing of viral spike protein priming for viral infection, and the cell surface protease transmembrane protease serine 2 (TMPRSS2) is highly expressed in the small airway epithelium^[4]. Since the outbreak of novel coronavirus, causing severe pneumonia and sequelae in late 2019, drug candidates for SARS-CoV have been re-tested for SARS-CoV-2. Wang et al. reported fruits tannic acids are dual inhibitors targeting viral main protease (3CL^{pro}) and host TMPRSS2 with IC₅₀ of 13.4 μM and 2.31 μM respectively^[5]. However, the potency is weak, and it appears that crude tannic acids cannot reach the therapeutic level without administration in large quantities.

Shih et al. further examined the antiviral activities of proprietary processed tannic acid from Chinese gallnuts, Ultrapure and Potent Tannic Acid (UPPTA). The EC₅₀ against SARS-CoV-2 is much improved to 0.5 μM in 3CL^{pro}-overexpressed cellular systems. Moreover, for the first documented human coronavirus infection, HCoV-OC43, the EC₅₀ is 1.3 μM. The selectivity index (EC₅₀/CC₅₀) declared its effectiveness and safety in vitro. They also evaluated the pharmacokinetics and toxicology of UPPTA to prove its feasibility in drug development^[6]. Also, a preliminary report reveals that patients infected by SARS-CoV-2 have a better clinical improvement^[7].

As SARS-CoV-2 directly targets the lungs and causes lung damage, treatment of COVID-19 with inhalants may offer more advantages over oral administration. Because the viruses incubate deep in the respiratory tract, the systemically administered drug could not reach them easily. Therefore, high doses of drugs are required through oral or parenteral routes and may cause systemic overexposure and unwanted side effects. The inhaled route of drug administration could provide a much higher concentration in the pulmonary system without much systemic exposure. Moreover, the peripheral metabolism of the drug could be greatly reduced when the drug is directly delivered to the lungs, minimizing systemic drug exposure^[8].

Since the toxicity of UPPTA by inhalation has never been investigated, a thorough evaluation is needed before human use and investigational new drug applications. In this study, we evaluated the potential systemic and pulmonary toxicity of UPPTA by inhalation in both Sprague-Dawley (SD) rats and Beagle dogs once a day for 2 consecutive weeks. We also investigated the reversibility of toxicity following a 3-week recovery period after the exposure.

2. Materials and methods

The purification and manufacture of UPPTA active pharmaceutical ingredient (API) was described in the previous research^[6]; the purity of cGMP-grade API was determined to be 98.4%.

2.1. Two-week toxicity study in SD rats following a 3-week recovery period

2.1.1. Aerosol generation and analysis

Inhalation exposure system and automatic metered-dose inhaler (MDI) nebulizer system (Beijing Huironghe Technology) were applied. Two bottles of control or test article were triggered to deliver the aerosol at the same time at the frequency of 12 times/min in single-cavity mode. The aerosol flow rate was set at 16 L/min, the dilution flow rate was 0 L/min, and the inner chamber exhaust flow rate is set at 14 L/min.

T_{99} of the exposure system was calculated by $4.6 \times V/a$, where V is the volume of the chamber (0.00321 m^3) and “ a ” (m^3/min) is the total aerosol in flow. During the experiment, the temperature was set at $20\text{--}26 \text{ }^\circ\text{C}$, the humidity was set at $30\text{--}80\%$, the oxygen concentration was set at $\geq 19\%$, and the carbon dioxide concentration was set at $< 1\%$.

For analysis of the concentration of UPPTA, samples were collected with a 3-stage wave plate sampler (5 mL of 30% acetonitrile water was added to each 1-stage wave tube) about 1 min after the beginning of administration. The sampling flow was set at 0.5 L/min. For analysis of the aerosol particle size parameters, including median mass aerodynamic diameter (MMAD), geometric standard deviation (GSD), and fine particle fraction, NGI laryngeal tube (NGI-1342 or NGI-1872, Copley Science, UK) and Copley Inhaler Testing Data Analysis Software, version 3.10 WIBU, were applied. Sample conditions were set at pumping flow rate: 15 L/min; collecting time: 3 min.

2.1.2. Animals and treatment

The study procedures were approved by the Institutional Animal Care and Use Committee. 8–9-week-old specific pathogen-free SD rats were purchased from the Zhejiang Vital River Laboratory Animal Technology Co. Ltd., China (license No. SCXK (Zhe) 2019-0001), housed at a maximum of 5 rats per cage and maintained on a 12/12-hour light/dark cycle at $23 \pm 4 \text{ }^\circ\text{C}$ and $55 \pm 15\%$ humidity. Animals had free access to SPF Rat Growth Breeding Feed diet (Jiangsu Xietong Pharmaceutical Bio-Engineering) and filtered water except during designated procedures. After the initial health examination, all animals were placed in a fixator for about 60 minutes once daily for 5 consecutive days for acclimation. Animals without abnormal behaviors during fixation acclimation were grouped and acclimated for an additional 5 days. All experiments were conducted in JOINN Laboratories (Suzhou) Co., Ltd. in accordance with the Good Laboratory Practice (GLP) guideline.

2.1.3. Experimental design

The objectives of this study are to evaluate the toxicity and toxicokinetic properties of UPPTA administered by MDI followed by a 3-week recovery period.

In the toxicity study, a total of 120 rats (60 rats/sex) were randomly assigned into 4 groups (15 animals/sex/group). Animals in Group 1 were given a 60-minute placebo by MDI as an excipient control. Animals in Groups 2 to 4 were given 15-minute, 30-minute, and 60-minute inhalations of UPPTA aerosol by MDI as low, medium, and high dose groups, respectively. The dosing frequency was once daily for 14 consecutive days for a total of 14 doses. The first dosing day was defined as Day 1. The actual delivered dose was calculated with the formula: dose (mg/kg) = ventilation volume per minute (RMV) \times concentration \times duration/BW/1000, whereas the RMV = $0.608 \times \text{BW (kg)}^{0.852}$.

Clinical observation (including daily clinical observation, detailed clinical observation, and local observation), body weight, food consumption, body temperature, ophthalmoscopic examinations, respiratory function (tidal volume, respiratory frequency, and minute respiratory volume), and clinical pathology (hematology, coagulation, clinical chemistry, and urinalysis) were evaluated during the study. Ten animals per sex per group were euthanized at the end of the dosing period (Day 15), and five animals per sex were euthanized at the end of the recovery period (Day 36) for macroscopic examination and complete necropsy examination in each group. Major organs were weighed, and histopathological examination was performed.

In the toxicokinetic study, a total of 56 rats (8 rats/sex of dosing groups; 4 rats/sex of control group) were randomly assigned into 4 groups. Animals in Group 1 were given a placebo by MDI for 60 min as

an excipient control. Animals in Groups 2 to 4 were given 15-minute, 30-minute, and 60-minute UPPTA aerosols by MDI as low, medium, and high dose groups, respectively. The dosing frequency was once daily for 14 consecutive days for a total of 14 doses. The first dosing day was defined as Day 1. The actual delivered dose was calculated with the formula: dose (mg/kg) = ventilation volume per minute (RMV) × concentration × duration/BW/1000, whereas the RMV = 0.608 × BW (kg)^{0.852}. Blood samples (approximately 0.3 mL, with EDTA-K₂ anticoagulation) were collected via jugular veins from animals of Groups 1 to 4 for toxicokinetic analysis.

2.1.4. Clinical observation

All rats were observed twice daily (a.m. and p.m.) during the study for signs of, but not limited to, mortality, moribundity, mental/behavior status, respiration, secretion, feces, and water and food intake.

Detailed clinical observations included, but were not limited to, mental status, behavior, skin, fur, eyes, ears, nose, abdomen, external genitalia, anus, limbs, feet, and respiration were conducted prior to first dosing and once weekly after dosing.

Local observation of irritation at the snout and other contact sites with aerosol and the reaction symptoms (such as swelling, asthma, cough, asphyxia, and other symptoms) of animals after dosing were conducted as well.

2.1.5. Clinical pathology

Blood samples for hematology, coagulation, and clinical chemistry determinations were collected via abdominal aorta prior to scheduled euthanasia. Rats were fasted overnight, then blood samples were collected into tubes containing EDTA-K₂, sodium citrate, or separation gel and coagulation promoters for hematology, coagulation, or clinical chemistry, respectively. For analysis, Sysmex[®] XN Hematology System, Sysmex[®] CS-2000i, Sysmex CS-5100, and TBA-120FR were used for hematology, coagulation, and clinical chemistry analysis, respectively. Urine was collected using metabolism cages and analyzed by the Cobas[®] 6500 automatic urine analyzer.

2.1.6. Toxicokinetic study

Blood samples were collected from animals of the control group at the pre-dose time point and 4 h after the end of dosing on Days 1 and 14. Blood samples were collected from animals of the dosing groups at the pre-dose time point, the end of dosing immediately (± 2 min), and at post-dosing at 30 min, 1 h, 2 h, 4 h, 8 h, and 24 h on Day 1 and Day 14. After blood collection, the blood samples were mixed by inverting at least 5 times and stored in an ice box. Plasma was separated by centrifugation at 2000 g for 10 min at 4 °C within 1 h after blood collection. The plasma was dispensed in duplicate into labeled centrifuge tubes and stored below -60 °C. The analysis of the biological sample was performed by a validated LC-MS/MS method for the determination of plasma levels.

2.1.7. Animal euthanasia and pathology examination

Animals were euthanized with isoflurane inhalation followed by abdominal aorta exsanguination on Day 15 and Day 36. For histopathological evaluation, tissues/organs were fixed in 10% neutral buffered formalin, followed by paraffin sectioning and hematoxylin and eosin (H&E) staining. A five-step grading system (minimal, slight, moderate, marked, or severe) was applied to rate the severity of microscopic lesions.

2.2. Two-week toxicity study in Beagle dogs following a 3-week recovery period

2.2.1. Aerosol generation and analysis

If not specified, the aerosol and its analysis were the same as those in the rodent study. For the dog experiment, two bottles of control or test article were triggered at the same time at the frequency of 8 times/min in single-cavity mode. The aerosol flow rate was set at 20 L/min, the dilution flow rate was 8 L/min, and the exhaust flow of the inner chamber was set at 26 L/min.

2.2.2. Animals and treatment

6 to 7-month-old conventional-grade Beagle dogs were purchased from Jiangsu Marshall Biotechnology (license No. SCXK (Zhe) 2019-0012), housed individually in stainless steel cages, and maintained on an approximate 12/12-hour light/dark cycle at 22 ± 4 °C and $55 \pm 15\%$ humidity. Animals had free access to a certified canine diet (Beijing Keaoxieli Feed) and tap water. After initial health examination, all animals underwent simulated domestication of inhalation for at least 5 days, about 135 minutes per day. Animals without abnormal behaviors during acclimation were grouped and acclimated for an additional 5 days. The study procedures were approved by the Institutional Animal Care and Use Committee.

2.2.3. Experimental design

To evaluate the potential toxicity and toxicokinetic properties, UPPTA was administered by MDI in Beagle dogs once a day for 2 consecutive weeks, and the reversibility of toxicity following a 3-week recovery period was also investigated.

A total of 40 Beagle dogs (20/sex) were randomly assigned into 4 groups (5/sex/group). Animals in Group 1 were given a 90-minute inhalation of placebo by MDI. Animals in Groups 2 to 4 were given 10-minute, 30-minute, and 90-minute UPPTA by MDI at low, medium, and high doses, respectively. Dosing frequency was once daily for 2 consecutive weeks for a total of 14 doses. The day of the first dosing was defined as Day 1.

Placebo group received excipient control. The actual delivered dose was calculated with the same formula as in the rodent study. Animals were evaluated for clinical observation, body weight, food consumption, body temperature, electrocardiogram, blood pressure, ophthalmological examination, and clinical pathology, which are similar to the rodent study. Blood samples were collected on Day 1 and Day 14 for toxicokinetic analysis. At the end of the dosing period, the first 3 animals/sex in Groups 1 to 4 were euthanized 2 weeks after dosing (Day 15). The remaining animals were euthanized at the end of the 3-week recovery period (Day 36). Systematic necropsy was performed for animals in Groups 1 to 4. Also, organ weight, gross anatomy, and histopathology were examined.

2.2.4. Clinical observation and clinical pathology

The clinical observation processes are the same as in the rat study. For clinical pathology, the equipment and procedures in the canine study are the same as those used in the rodent study except for the ADVIA 2120 Hematology System, which was used for hematology, and the blood samples were collected via forelimb vein.

2.2.5. Toxicokinetic study

Blood samples (approximately 1 mL) were collected via forelimb subcutaneous vein for bioanalysis and toxicokinetic analysis. Blood samples were collected from animals of the control group at the pre-dose time point and at post-dosing 2 h on Day 1 and Day 14. Blood samples were collected from each animal in the dosing groups at the pre-dose time point, after the end of dosing immediately, and at post-

dosing 2, 4, 8, 12, and 23 h on Day 1. On Day 14, blood samples from 3/sex animals of the dosing groups were collected at the pre-dose time point, at the end of dosing immediately, and at post-dosing 2, 4, 8, 12, and 23 h. Blood samples from other 2/sex animals of dosing groups were collected at the pre-dose time point, at the end of dosing immediately, and at post-dosing 2, 4, 8, 12, 23, 48, 72, 96, 120, 144, 264, 384, and 504 h.

2.2.6. Animal euthanasia and pathology examinations

Animals were euthanized after intramuscular injection of Xoletil 50 (12 mg/kg, 50 mg/mL) and xylazine hydrochloride (1 mg/kg, 20 mg/mL), followed by femoral artery bloodletting on Day 15 and Day 36. Histopathological evaluations were the same as in the rodent study.

2.3. Test article concentration in lung tissue

UPPTA was administered by MDI in SD rats and Beagle dogs once a day for 1 consecutive week. Frozen rats and dog lung samples were completely thawed in a cooler filled with ice and homogenized well. 20 μ L rat lung matrix, 50 μ L of internal standard (IS) (200 ng/mL), and 300 μ L of tannase reaction solution (1 mg/mL) were added into an Eppendorf tube and mixed well by vortex. The mixtures were incubated at 30 °C for 4 h to complete the hydrolysis reaction. The hydrolysate was extracted with 3300 μ L of extraction reagent (1.5% (w/w) formic acid in ACN), vortexed for about 2 min and then centrifuged at 12,000 rpm for 10 min at 2–8 °C. The supernatant was collected and followed by evaporating to dryness under N₂. The dried extract was reconstituted in 200 μ L reconstitution solution and filtered through a 0.22 μ m membrane filter. The analysis of the biological sample was performed by a validated LC-MS/MS method for the determination of tissue levels.

2.4. Statistical analysis

Data from male and female animals were analyzed separately. All statistical tests were conducted as 2-sided tests, and the level of significance was set at 0.05 or $p \leq 0.05$. Group means and standard deviations (mean \pm SD) were calculated using the Provantis system (SAS 9.2). Data were analyzed with the Levene's test to assess variance homogeneity. Data with homogeneous variance ($p > 0.05$ in Levene's test) were analyzed with the one-way ANOVA followed by the Dunnett's test for multiple comparisons. Data with variance heterogeneity ($p \leq 0.05$ in Levene's test) were logarithmically transformed using the natural logarithm (ln transformation). The Kruskal-Wallis test (non-parametric method) followed by the two-independent-sample test (Mann-Whitney U) for multiple comparisons was performed on the original data when transformed data showed variance heterogeneity.

When a dataset contained negative values, the logarithmical transformation was not applied, and either the original data or the rank-transformed data were used for statistical analysis. When a dataset had zero values, the zero values were regarded as 1/10 of the smallest positive value in the dataset when the logarithmical transformation was performed.

3. Results

3.1. Analysis findings of the test article

In the SD rodent study, aerosol concentrations were analyzed from Day 1 to 14, and the aerosol particle size distribution was analyzed after the dosing on Days 1, 7, and 14. The actual delivered dose was 2.88 ± 0.96 mg/kg for the low-dose group, 5.35 ± 1.28 mg/kg for the medium-dose group, and 11.70 ± 2.20 mg/kg for the high-dose group. The actual delivered dose and aerosol particle size distribution parameters are presented in **Table 1**.

Table 1. The actual delivered dose and aerosol particle size distribution parameters.

Dose	Low	Medium	High
SD rat			
Target dose (mg/kg)	3	6	12
Actual delivered dose (mg/kg)	2.88 ± 0.96	5.35 ± 1.28	11.70 ± 2.20
Average MMAD (µm)	3.11	3.07	3.26
Average GSD	1.55	1.56	1.57
Average FPF (%)	85.49	85.33	81.1
T ₉₉ (min)	0.92		
Beagle dog			
Target dose(mg/kg)	0.3	1.0	3.0
Actual delivered dose(mg/kg)	0.28 ± 0.09	0.98 ± 0.18	3.05 ± 0.48
Average MMAD(µm)	3.37	3.37	3.37
Average GSD	1.49	1.49	1.49
Average FPF (%)	82.81%	82.81%	82.81%
T ₉₉ (min)		0.49	

Note: MMAD: Mass Median Aerodynamic Diameter, GSD: Geometric Standard Deviation, FPF: Fine Particle Fraction (Fine Particle Fraction, MMAD < 5 µm). The T₉₉ calculation formula of the exposure system is $T_{99} = 4.6 \times V/a$, V is the exposure tower volume (0.00321 m³), a is the set total aerosol generation and dilution flow (m³/min), a = 0.016 m³/min for this test.

In the Beagle dog study, aerosol concentrations were analyzed from Days 1 to 14, and the aerosol particle size distribution was analyzed after the dosing on Days 1, 7, and 14. The actual delivered dose was 0.28 ± 0.09 mg/kg in the low-dose group, 0.98 ± 0.18 mg/kg in the medium-dose group and 3.05 ± 0.48 mg/kg in the high-dose group, respectively. The T₉₉ (min) of the exposure system was 0.49 min ($T_{99} = 4.6 \times V/a$ (V = 0.003 m³, a = 0.028 m³/min)). The actual delivered dose and aerosol particle size distribution parameters are in **Table 1**.

3.2. Clinical observation and clinical pathology

In the rodent study, there are no test article-related abnormal changes (including daily clinical observation, detailed clinical observation, and local observation), body weight, food consumption, body temperature, ophthalmoscopic examinations, respiratory function, and clinical pathology (hematology, coagulation, clinical chemistry, and urinalysis) in all the dose groups during the study.

In the Beagle dog study, no mortality related to the test article was observed in all three dose groups during the whole study. There are no abnormalities related to the test article, and no toxicologically significant changes were observed in body weight, food consumption, body temperature, electrocardiogram (limb II ECG), blood pressure, ophthalmic examination, coagulation, clinical chemistry, urinalysis, and gross anatomy in all the animals.

3.3. Toxicokinetic study

In the SD rodent study, Airneclu[®], UPPTA was not detected after 15 min, 30 min, and 60 min inhalation in most of the plasma samples.

In the Beagle dog study, after the first (Day 1) and last dosing (Day 14), Airneclu[®], UPPTA was not detected in all the plasma samples of the excipient control group. The key TK parameters of Airneclu[®] and UPPTA in the low, middle, and high dose groups are shown in **Table 2**.

Table 2. The key toxicokinetic parameters of Airnecflu®, UPPTA in the Beagle dog study.

Analyte	Study day	Dose level (mg/kg)	Gender	T _{max} (hr)	C _{max} (ng/mL)	AUC ₀₋₂₃ (hr·ng/mL)
Airnecflu®	D1 #	0.28	Male	-	-	-
			Female	-	-	-
UPPTA	0.98	0.98	Male	15.67 ± 6.35	103.02 ± 49.40	1453.46 ± 719.50
			Female	12.00 ± 0.00	81.93 ± 36.90	1233.66 ± 739.20
	3.05	3.05	Male	14.20 ± 4.92	230.93 ± 98.45	3920.88 ± 1593.09
			Female	12.00 ± 0.00	356.43 ± 129.60	6237.47 ± 1984.03
	D14 #	0.28	Male	0.40 ± 0.89	90.10 ± 21.52	1773.31 ± 338.18
			Female	0.80 ± 1.79	126.14 ± 63.62	2785.10 ± 1219.51
0.98	0.98	Male	2.40 ± 3.57	432.62 ± 121.85	8621.89 ± 2491.84	
		Female	1.61 ± 3.57	738.87 ± 475.47	15077.32 ± 10791.66	
3.05	3.05	Male	2.01 ± 1.99	1162.13 ± 613.63	24206.36 ± 12767.67	
		Female	7.21 ± 6.56	1684.23 ± 841.22	34553.42 ± 16559.60	

∴ Data under detectable limits; #, *n* = 5 for each group.

The analysis of the two-sided *t*-test was applied in comparing the sexual difference of toxicokinetic parameters. The result showed that the exposure (AUC₀₋₂₃) of Airnecflu®, UPPTA, had no statistical difference (*p* > 0.05) in both genders after inhalation of Airnecflu®, UPPTA at doses of 0.28, 0.98, and 3.05 mg/kg. The ratios (female/male) of mean C_{max} ranged from 0.80 to 1.71, and the ratios (female/male) of mean AUC₀₋₂₃ ranged from 0.85 to 1.75 on the first dosing day (D1) and after repeated administration (D14). No gender differences were observed in the exposures of Airnecflu®, UPPTA.

In the Beagle dogs, after inhalation of Airnecflu®, UPPTA aerosol at doses of 0.98 and 3.05 mg/kg on Day 1, the mean exposure (AUC₀₋₂₃) in plasma increased dose-dependently in both genders. On Day 14, after inhalation of Airnecflu®, UPPTA aerosol at doses of 0.28, 0.98, and 3.05 mg/kg for 2 consecutive weeks, the mean exposure (AUC₀₋₂₃) increased dose-dependently in both genders. The ratio of mean exposure was slightly greater than the ratio of dose when comparing low dose to medium or high dose.

After consecutive 14-day administration of Airnecflu®, UPPTA at 0.98 and 3.05 mg/kg levels to Beagle dogs, the mean of Airnecflu®, UPPTA C_{max} ratios (Day14/Day1) ranged from 5.53 to 9.84 and the mean of AUC₀₋₂₃ ratios (Day14/Day1) ranged from 6.78 to 13.87 (Table 2).

3.4. Organ weight change and pathology

In the SD rat study, test article-related increases in the absolute organ weight, organ-to-body weight ratio, and organ-to-brain weight ratio of lung with bronchi were observed in all dosing groups at the terminal necropsy (Day 15) (Table 3). These organ weight changes were probably correlated to the following pathology changes.

In the SD rats, test article-related minimal to slight alveolar macrophage aggregation and minimal to moderate perivascular/interstitial inflammatory cell infiltration were noted in the lung with bronchi of both sexes. Minimal alveolar epithelium hyperplasia was noted only in the high-dose group in the lung with bronchi (Figure 1 and Table 4). After a 3-week recovery period (Day 36), there is substantial recovery of the test article-related microscopic findings in the lung with bronchi.

Table 3. Weight changes in terminal necropsy and their recover on Day 36 of the SD rats.

Sex	Male			Female		
Dose	Low	Medium	High	Low	Medium	High
Inhalation duration (min)	15	30	60	15	30	60
Terminal necropsy (Day 15) #						
Lung with bronchi						
Organ weight	1.61 g*	1.56 g*	1.69 g*	1.29 g	1.32 g*	1.34 g*
Organ-to-Body	0.46%*	0.46%*	0.48%*	0.52%*	0.52%*	0.55%*
Organ-to-Brain	78.47%*	78.28%	85.76%*	67.35%	67.46%	71.00%
Recovery necropsy (Day 36) ^						
Lung with bronchi						
Organ weight	1.53 g	1.59 g	1.67 g	1.27 g	1.23 g	1.24 g
Organ-to-Body	0.35%	0.35%	0.38%	0.47%	0.47%	0.47%
Organ-to-Brain	74.42%	78.70%	80.65%	64.00%	62.88%	65.06%

Data Format: Control data were presented in absolute (actual) values for the control group only; the data for dosed animal were presented as a percentage of change relative to the control group data. * $p \leq 0.05$; #, $n = 10$ for each group; ^, $n = 5$ for each group.




Figure 1. Examples of test article-related moderate alveolar macrophage aggregation, moderate interstitial/alveolar inflammatory cells infiltration, and minimal alveolar epithelium hyperplasia were noted in the lung with bronchi in SD rat. Representative lung H&E staining sections of rats with various treatments were displayed. Scale bar = 100 µm.

Table 4. UPPTA-related microscopic findings at the terminal and recovery necropsy in SD rat study.

Sex	Male				Female			
Dose	Control	Low	Medium	High	Control	Low	Medium	High
Inhalation duration/time (min)	60	15	30	60	60	15	30	60
Terminal necropsy (Day 15) #								
Lung with bronchi								
Alveolar macrophage aggregation								
Minimal	0	8	6	3	0	1	6	3
Slight	0	0	0	6	0	0	0	6
Infiltration, inflammatory cells perivascular; interstitial								
Minimal	0	5	3	1	0	10	6	1
Slight	0	4	5	2	0	0	4	3
Moderate	0	0	0	7	0	0	0	6
Hyperplasia, alveolar epithelium								
Minimal	0	0	0	1	0	0	0	2
Recovery necropsy (Day 36) *								
Lung with bronchi								
Alveolar macrophage aggregation								
Minimal	0	2	0	2	0	0	1	0
Infiltration, inflammatory cells perivascular; interstitial								
Minimal	0	2	2	2	0	0	1	3

#, $n = 10$ for each group; *, $n = 5$ for each group.

In the Beagle dog study, varying degrees of test article-related increases in the absolute weight, organ-to-body weight ratio, and organ-to-brain weight ratio of the lung with bronchi were observed in all dosing groups at the terminal necropsy (Day 15) (**Table 5**). These organ weight changes were probably correlated to the following pathology changes.

Table 5. Weight changes in terminal necropsy and their recover on Day 36 of the Beagle dog.

Sex	Male			Female		
Dose	Low	Medium	High	Low	Medium	High
Inhalation duration (min)	10	30	90	10	30	90
Terminal necropsy (Day 15) #						
Lung with bronchi						
Organ weight	72.12 g	87.26 g*	87.69 g*	64.63 g	71.08 g	78.32 g
Organ-to-Body	0.86%	1.01%	1.05%*	0.98%	1.08%	1.18%
Organ-to-Brain	100.97%	118.09%	122.51%	90.24%	106.19%	112.93%
Recovery necropsy (Day 36) ^						
Lung with bronchi						
Organ weight	73.00 g	72.50 g	76.15 g	68.17 g	61.60 g	79.88 g
Organ-to-Body	0.88%	0.88%	0.93%	0.96%	0.87%	1.05%
Organ-to-Brain	105.53%	93.72%	97.50%	96.33%	90.18%	111.99%

Note: Data Format: Control data is presented in absolute terms; the data for dosed animals is presented as a percentage of change relative to the concurrent control group data. * $p \leq 0.05$, #, $n = 3$ for each group; ^, $n = 2$ for each group.

There was test article-related minimal to moderate alveolar macrophage aggregation, minimal to moderate perivascular/interstitial inflammatory cell infiltration, and minimal alveolar epithelium hyperplasia in both sexes' lungs with bronchi. In addition, test article-related minimal to slight increased cellularity of lymphocytes in cortex and minimal inflammatory cells infiltrated in medulla were noted in both sexes' bronchial lymph nodes (**Table 6**). After a 3-week recovery period (Day 36), although test article-related microscopic findings were still observed in the lung with bronchi, the severity and incidence of the above lesions were significantly lower than those observed at the terminal necropsy (Day 15), indicating a substantial recovery.

Table 6. UPPTA-related microscopic findings at the terminal and recovery necropsy in the Beagle dog study.

Sex	Male				Female			
	Control	Low	Medium	High	Control	Low	Medium	High
Dose	Control	Low	Medium	High	Control	Low	Medium	High
Inhalation duration (min)	90	10	30	90	90	10	30	90
Terminal necropsy (Day 15) #								
Lung with bronchi								
Alveolar macrophage aggregation								
Minimal	0	0	1	0	0	0	2	0
Slight	0	0	1	2	0	0	1	2
Moderate	0	0	0	1	0	0	0	1
Infiltration, Inflammatory cells, interstitium/perivascular/alveous								
Minimal	1	0	1	0	1	2	1	0
Slight	0	0	1	2	0	0	2	2
Moderate	0	0	0	1	0	0	0	1
Hyperplasia, alveolar epithelium, bronchial epithelium								
Minimal	0	0	1	3	0	0	0	1
Lymph node, bronchial								
Increased cellularity, lymphocytic, cortex								
Minimal	0	1	0	1	0	0	2	0
Slight	0	0	1	2	0	0	1	3
Infiltration, inflammatory cells, medulla								
Minimal	0	0	1	0	0	0	1	3
Recovery necropsy (Day 36) *								
Lung with bronchi								
Infiltration, inflammatory cells, interstitium/perivascular								
Minimal	0	1	2	0	0	1	0	1

#, n = 3 for each group; *, n = 2 for each group.

3.5. Test article concentration in lung tissues

Airneclu® UPPTA aerosol was administered to SD rats and Beagle dogs by repeated inhalation at actual mean delivered doses of 26.87 mg/kg and 8.10 mg/kg, respectively, once daily for 7 consecutive days. Test article concentration was measured in lung tissues after animal euthanasia. 425.90 and 10.46 µg UPPTA per gram tissue were detected in 3 rat whole lung tissues and 2 canine accessory lobes of lung, respectively.

4. Discussion

Tannic acids have been used in many regions worldwide for multiple therapeutic applications. In Eastern countries, they were applied in traditional treatments in the herbal recipe for various diseases such as diarrhea, bleeding, toxification, dementia, and coughing^[1,9,10]. In Western countries, tannic acids were used for toxification, burn injuries, and diarrhea^[11–13].

Commercial crude tannic acids are derived from various origins, such as sumac galls, Aleppo oak galls, or sumac leaves. Among these, Chinese gallnut (*Galla chinensis*), which is produced by the Chinese aphid (*Schlechtendalia chinensis*) through stimulation of sumac leaves (*Rhus chinensis*), is the main resource of tannic acids in East Asia for traditional Chinese medicine^[14,15].

The coronavirus' lifecycle starts with binding of the envelope spike protein to its receptor, angiotensin-converting enzyme 2 (ACE2). Through member fusion, the RNA genome is released into the host cytosol, where it is translated into the replicase proteins. The polyproteins, pp1a and pp1ab, are cleaved by a virus-encoded protease (3CL^{pro}) into individual replicase complex nonstructural proteins. Replication begins in virus-induced double-membrane vesicles derived from the endoplasmic reticulum (ER). The positive-strand genome then serves as a template for full-length negative-strand RNA and sub-genomic RNA (sgRNA). sgRNA translation results in the synthesis of both structural proteins and accessory proteins that are inserted into the ER-Golgi intermediate compartment (ERGIC) for virion assembly. Finally, subsequent positive-sense RNA genomes are incorporated into newly synthesized virions, which are then secreted from the cell membrane^[16–18].

Because of the key role of 3CL^{pro} in proteolytic processing during coronavirus replication, it has been considered a critical molecular target for the development of anti-coronavirus drugs. In 2005, Chen et al. screened 720 natural compounds for inhibitory activity against SARS-CoV. Tannic acids, extracted from tea, showed the highest inhibitory activity against 3CL^{pro} with IC₅₀ of 3 μM^[3]. In late 2019, a novel coronavirus causing a severe pneumonia outbreak was lately identified and named SARS-CoV-2. Drug candidates once tested for SARS-CoV were re-tested for being potential treatments for SARS-CoV-2.

Wang et al. reported a molecular analysis that demonstrated that tannic acids from fruit formed a thermodynamically stable complex with SARS-CoV-2 3CL^{pro} at a KD of 1.1 μM^[5]. They also tested the inhibitory activity of tannic acid against SARS-CoV-2 3CL^{pro} with IC₅₀ of 13.4 μM. 3CL^{pro} of SARS-CoV and SARS-CoV-2 are 96% identical at the amino acid level, and only one different residue is presented at the binding pocket^[19,20]. It is therefore predicted that tannic acids are effective in inhibiting both SARS-CoV-2 activity in addition to SARS-CoV.

Shih et al. further tested the activities of proprietary purified tannic acids from Chinese gallnut, named ultrapure and potent tannic acid (UPPTA), Pentarlandir[®]^[6]. The EC₅₀ against two coronaviral strains, SARS-CoV-2 and HCoV-OC43, the earliest-known human coronavirus, are 0.5 μM and 1.3 μM, respectively, as tested in the 3CL^{pro}-overexpressed cellular system. Even though SARS-CoV-2 shows only 59% identity of 3CL^{pro} at the amino acid level with HCoV-OC43 in the binding pocket^[20], the inhibitory activities of Pentarlandir[®] are very close in both strains. It declares the potential of proprietary tannic acids as broad-spectrum pancoronavirus drugs to treat other human coronaviruses with similar 3CL^{pro} amino acid identities, such as HCoV-OC43.

The UPPTA is a tannic acid derived from Chinese gallnut (*Galla chinensis*) with a high proportion of 5–12 galloyl moieties as compared to other sources of tannic acid from *Quercus infectoria* and *Rhus chinensis*. In fact, the inhibitory activities of tannic acids are positively correlated with the number of

galloyl groups^[21]. In addition, the selectivity index (EC_{50}/CC_{50}) of the coronaviruses is above 100 when treating UPPTA, which declares its effectiveness and safety. The pharmacokinetic (PK) and toxicology of UPPTA were also investigated by oral administration. In the PK studies, proprietary processed tannic acids were distributable to the pulmonary system without accumulation^[6]. A 14-day repeated dose toxicity study in rats demonstrated its safety with a NOAEL (no-observed-adverse-effect level) value above 2000 mg/kg/day^[6]. These findings support UPPTA for further drug development for coronaviruses.

Though the safety profile of UPPTA by oral administration has been examined thoroughly, the potential toxicity of Airneclu[®] UPPTA by MDI administration has never been investigated. In this study, we evaluated the potential toxicity and the target organ toxicity of Airneclu[®] UPPTA by inhalation in both SD rats and Beagle dogs, once a day for 2 consecutive weeks. All the procedures were under GLP regulations.

In our previous studies, we evaluated the acute and 7-day repeated dose toxicity of Airneclu[®] UPPTA by inhalation in SD rats and Beagle dogs. The actual mean delivered doses used in the single dose and 7-day repeated dose toxicity of the rat studies were up to 139.20 and 26.87 mg/kg/day, respectively. In the Beagle dog studies, the actual mean delivered doses used in single dose and 7-day repeated dose toxicity were up to 12.40 and 8.10 mg/kg/day, respectively. These doses were defined as maximum tolerated dose (MTD) as there was no death, mortality, and/or significant systemic toxicity observed. Therefore, the expected delivered doses in the 14-day repeatedly toxicological evaluation were set at 12.00 and 3.00 mg/kg/day in SD rat and Beagle dog studies, respectively.

In this study, our findings revealed no death, mortality, or significant systemic toxicity at the actual mean delivered dose of Airneclu[®] UPPTA up to 11.70 ± 2.20 mg/kg and 3.05 ± 0.48 mg/kg in SD rats and Beagle dogs, respectively. In the pathology study, test article-related changes, including alveolar macrophage aggregation, perivascular/interstitial inflammatory cell infiltration, and alveolar epithelium hyperplasia, were mostly minimal to slight in SD rats and Beagle dogs. Furthermore, after a 3-week period, substantial recovery of the test article-related microscopic findings was observed in the lung with bronchi in both SD rats and Beagle dogs, indicating a reversible pathology that can recover.

In the toxicokinetic study, Airneclu[®] UPPTA was not detected in most of the plasma samples of rats, indicating that it rarely enters the peripheral circulation after inhalation. However, in the canine toxicokinetic study, Airneclu[®] UPPTA was detected in the plasma of a Beagle dog at a very low level ($C_{max} \sim 1$ μ M; molecular weight of UPPTA is 1465 g/mol), and the mean exposure (AUC_{0-23}) increased dose-dependently. The ratio of mean exposure was slightly greater than the ratio of dose when comparing low dose to medium or high dose after 14 days of administration. We also measured the concentration of Airneclu[®] UPPTA in lung tissues at the end of the 7-day dosing period from the previous study. Substantial levels of UPPTA were detected in the lung tissues. Taken together, there is a significant gradient of distribution of UPPTA when administered by MDI; UPPTA can be administered successfully by MDI to pulmonary tissue, and there is little UPPTA distributed systemically.

Janower et al. reported potential hepatotoxicity caused by tannic acid through enema administration in a few cases after barium enema examinations^[22]. However, liver toxicity was considered to be secondary to tannic acid administration with the performance of barium enemas. Shih et al. conducted the 14-day repeated dose toxicity study by oral route under GLP regulations. They evaluated a full panel of toxicology endpoints and found that the purified tannic acid had no significant abnormalities in all organs, biochemical parameters, or other evaluations^[6]. In our phase II human study of UPPTA by oral

administration, no serious adverse event (SAE) was noted, and the adverse events (AE) were minimal and negligible^[7].

We have reported the virucidal activity of proprietary UPPTA against SARS-CoV-2. Moreover, we have also investigated the virucidal activity of UPPTA against influenza viruses. UPPTA also showed virucidal activity to inhibit the infection of H1N1 and H3N2 at sub- μ M concentrations in vitro. In addition, UPPTA further improved influenza-induced body survival rate, weight loss, and subsequent lung damage in vivo (unpublished findings).

Overall, the administration of UPPTA by either oral administration or MDI inhalation was well tolerated. The minimal pathology observed by MDI administration is, to a large extent, reversible. The no observed adverse effect levels (NOAEL) for MDI administration of UPPTA are 11.70 ± 2.20 mg/kg in SD rats and 3.05 ± 0.48 mg/kg in Beagle dogs, respectively. Airnecflu[®] UPPTA is considered safe for future use against either influenza or coronavirus infection, as it can be effectively delivered to reach therapeutic concentration in the target organ and minimize systemic exposure.

Author contributions

Conceptualization, LPL, YWM and GET; methodology, LPL, YWM and GET; software, LPL and YWM; validation, LPL and YWM; formal analysis, LPL and GET; investigation, LPL and TT; resources, TYC and GET; data curation, LPL, YWM and TT; writing—original draft preparation, LPL and YAL; writing—review and editing, LPL, YAL, TT and GET; visualization, LPL and GET; supervision, TYC and GET; project administration, LPL and YWM; funding acquisition, GET. All authors have read and agreed to the published version of the manuscript.

Funding

This study is supported and funded by SyneuRx International (Taiwan) Corporation. LPL, YWM, and YAL are employees of SyneuRx International (Taiwan) Corporation.

Conflict of interest

The authors declare no conflict of interest.

References

1. Tian F, Li B, Ji B, et al. Identification and structure–activity relationship of gallotannins separated from *Galla chinensis*. *Lebensmittel-Wissenschaft und-Technol* 2009; 42: 1289–1295.
2. Green RH. Inhibition of multiplication of influenza virus by tannic acid. *Experimental Biology and Medicine* 1948; 67(4): 483–484. doi: 10.3181/00379727-67-16347p
3. Chen CN, Lin CPC, Huang KK, et al. Inhibition of SARS-CoV 3C-like protease activity by theaflavin-3,3'-digallate (TF3). *Evidence-Based Complementary and Alternative Medicine* 2005; 2(2): 209–215. doi: 10.1093/ecam/neh081
4. Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 2020; 181(2): 271–280. doi: 10.1016/j.cell.2020.02.052
5. Wang SC, Chen Y, Wang YC, et al. Tannic acid suppresses SARS-CoV-2 as a dual inhibitor of the viral main protease and the cellular TMPRSS2 protease. *American Journal of Cancer Research* 2020; 10(12): 4538–4546.
6. Shih PC, Mao YW, Hu JW, et al. Development of ultrapure and potent tannic acids as a pan-coronal antiviral therapeutic. *ACS Pharmacology & Translational Science* 2022; 5(6): 400–412. doi: 10.1021/acspsci.1c00264
7. Wu V, Lu LP, Chang WH, et al. Pentarlandir[®] for the treatment of adult with early COVID-19: A randomized, double blind, placebo controlled, proof of principle study. Unpublished data.

8. Rau JL. The inhalation of drugs: Advantages and problems. *Respir Care* 2005; 50(3): 367–382.
9. Fatima S, Haider N, Alam A, et al. Preventive, promotive and curative aspects of dementia in complementary medicine (Unani): Through-Black Box Design. *International Journal of Herbal Medicine* 2017; 5(1): 1–5.
10. Djakpo O, Yao W. *Rhus chinensis* and *Galla Chinensis* – folklore to modern evidence: Review. *Phytotherapy Research* 2010; 24(12): 1739–1747. doi: 10.1002/ptr.3215
11. Daly JS, Cooney DO. Interference by tannic acid with the effectiveness of activated charcoal in “Universal Antidote.” *Clinical Toxicology* 1978; 12(5): 515–522. doi: 10.3109/15563657809150025
12. Chokotho L, van Hasselt E. The use of tannins in the local treatment of burn wounds—A pilot study. *Malawi Medical Journal* 2005; 17: 19–20. doi: 10.4314/mmj.v17i1.10866
13. Loeb H, Vandenplas Y, Würsch P, et al. Tannin-rich carob pod for the treatment of acute-onset diarrhea. *Journal of Pediatric Gastroenterology and Nutrition* 1989; 8(4): 480–485. doi: 10.1097/00005176-198905000-00010
14. Chen H, Liu J, Cui K, et al. Molecular mechanisms of tannin accumulation in *Rhus* galls and genes involved in plant-insect interactions. *Scientific Reports* 2018; 8(1). doi: 10.1038/s41598-018-28153-y
15. Ren Y, Zhang X, Li T, et al. *Galla chinensis*, a traditional Chinese medicine: Comprehensive review of botany, traditional uses, chemical composition, pharmacology and toxicology. *Journal of Ethnopharmacology* 2021; 278: 114247. doi: 10.1016/j.jep.2021.114247
16. Lee C. Porcine epidemic diarrhea virus: An emerging and re-emerging epizootic swine virus. *Virology Journal* 2015; 12(1). doi: 10.1186/s12985-015-0421-2
17. Jeong GU, Song H, Yoon GY, et al. Therapeutic strategies against COVID-19 and structural characterization of SARS-CoV-2: A review. *Frontiers in Microbiology* 2020; 11. doi: 10.3389/fmicb.2020.01723
18. Yadav R, Chaudhary JK, Jain N, et al. Role of structural and non-structural proteins and therapeutic targets of SARS-CoV-2 for COVID-19. *Cells* 2021; 10(4): 821. doi: 10.3390/cells10040821
19. Zhang L, Lin D, Sun X, et al. Crystal structure of SARS-CoV-2 main protease provides a basis for design of improved α -ketoamide inhibitors. *Science* 2020; 368(6489): 409–412. doi: 10.1126/science.abb3405
20. Zhang Y, Gao H, Hu X, et al. Structure-based discovery and structural basis of a novel broad-spectrum natural product against the main protease of coronavirus. *Journal of Virology* 2022; 96(1). doi: 10.1128/jvi.01253-21
21. Mao YW, Lu LP, Shih PC, et al. Ultrapure and potent tannic acid (UPPTA) is a novel inhibitor of D-amino acid oxidase to improve the N-methyl-D-aspartate function of CNS disorders. *Phytomedicine Plus* 2023; 3(1): 100399. doi: 10.1016/j.phyplu.2022.100399
22. Janower ML, Robbins LL, Tomchik FS, et al. Tannic Acid and the Barium Enema. *Radiology* 1965; 85(5): 887–894. doi: 10.1148/85.5.887